



Asahikawa Medical College Repository <http://amcor.asahikawa-med.ac.jp/>

Human Reproduction (2008) 23(2):233–239.

Mouse and human spermatozoa can be freeze-dried without damaging their chromosomes.

Kusakabe H, Yanagimachi R, Kamiguchi Y.

**Title:**

Mouse and human spermatozoa can be freeze-dried without damaging their chromosomes

5 **Running title:**

Freeze-dried mouse and human spermatozoa

**Authors:**

H.Kusakabe<sup>1,3</sup>, R.Yanagimachi<sup>2</sup> and Y.Kamiguchi<sup>1</sup>

10

<sup>1</sup> Department of Biological Sciences, Asahikawa Medical College, 2-1-1-1

Midorigaoka-higashi, Asahikawa 078-8510, Japan

<sup>2</sup> Institute for Biogenesis Research, University of Hawaii Medical School, 1960

East-West Road, Honolulu, Hawaii 96822, USA

15 <sup>3</sup> To whom correspondence should be addressed at: Hirokazu Kusakabe, Department  
of Biological Sciences, Asahikawa Medical College, 2-1-1-1 Midorigaoka-higashi,  
Asahikawa 078-8510, Japan. E-mail: hkusa55@asahikawa-med.ac.jp

**Key words: ICSI/freeze-drying/human sperm/mouse sperm/chromosome**

20

## Abstract

**BACKGROUND:** Although mouse spermatozoa can be freeze-dried without losing their reproductive capacity, the technique needs further improvements to reduce the incidence of chromosomal damages to spermatozoa. Effects of freeze-drying on human

spermatozoa are unknown. **METHODS:** Mouse spermatozoa were suspended in a Tris-buffered EGTA solution briefly (about 10 min at 37°C) or for 1-7 days at 4°C before freeze-drying. Freeze-dried spermatozoa were maintained for up to 1 year at 4°C before injection. Sperm chromosomes were examined during the first mitosis (cleavage)

of zygotes. Sperm's ability to support embryo development was assessed by examining mid-gestation fetuses after transfer of 2-cell embryos to surrogate mothers. Chromosome integrity of freeze-dried human spermatozoa was examined by injecting individual spermatozoa into mouse oocytes which were previously enucleated.

**RESULTS:** When mouse spermatozoa were freeze-dried immediately after suspension in Tris-buffered EGTA solution, only about 40% had normal chromosomes. When the spermatozoa were kept in the same solution for 3-7 days before freeze-drying, 85-95% had normal chromosomes and they were able to support embryo development better than those which were in the solution briefly. Freeze-dried human spermatozoa well maintained their chromosomes regardless of the duration of pre-freeze-drying

incubation of spermatozoa in the Tris-buffered EGTA solution. **CONCLUSIONS:** Prior incubation of mouse spermatozoa in Tris-buffered EGTA solution for several days makes sperm chromosomes more resistant to freeze-drying. As the consequence, spermatozoa freeze-dried this way support embryo development better than those exposed to Tris-buffered EGTA solution only briefly. Freeze-dried human spermatozoa

well maintained their chromosomes without pre-freeze-drying incubation in Tris-buffered EGTA solution.

## Introduction

5

Long-term preservation of mammalian spermatozoa is now possible by various methods including: (1) deep freezing with cryoprotection (Polge *et al.*, 1949; Nakagata 2000); (2) deep freezing without cryoprotection (Wakayama *et al.*, 1998a; Ward *et al.*, 2003); (3) freeze-drying (Wakayama and Yanagimachi 1998; Kusakabe *et al.*, 2001); (4) 10 drying without freezing (Bhowmick *et al.*, 2003; McGinnis *et al.*, 2005; Li *et al.*, 2007) and (5) freezing whole animal body or of isolated testes (Ogonuki *et al.*, 2006). In all but the first, spermatozoa must be injected microsurgically into oocytes because they are motionless or “dead” in the conventional sense. An advantage of freeze-drying or drying spermatozoa is obvious. Neither liquid nitrogen nor dry ice is needed for the 15 storage and shipment of preserved spermatozoa. Someday, it could be possible to store spermatozoa indefinitely at ambient temperature and ship them to any parts of the world without dry ice or liquid nitrogen.

Ward *et al.* (2003) compared chromosome integrity and reproductive potential of freeze-dried mouse spermatozoa with those frozen in liquid nitrogen without any 20 cryoprotection. When examined 12 months after storage at 4°C, the frozen spermatozoa stored in liquid nitrogen supported embryo development better than freeze-dried spermatozoa stored at 4°C. Structural chromosome aberrations were less common in frozen spermatozoa than in freeze-dried spermatozoa. We report here that chromosome damage in freeze-dried mouse spermatozoa can be alleviated when spermatozoa are

kept in an EGTA solution for several days (at 4°C) before freeze-drying. To our surprise, human spermatozoa do not require this extra step to maintain their chromosome integrity.

## 5      **Materials and Methods**

### *Animals*

Oocytes and spermatozoa were collected from hybrid mice (B6D2F<sub>1</sub>) at 7-12 weeks of age. Fully mature albino females [Crlj:CD-1(ICR)] were used as surrogate mothers  
10 after mating with vasectomized albino males [Crlj:CD-1(ICR)] the day before embryo transfer. The females were maintained under 14-h light/10-h dark cycles at 22-24°C. Food and water were given *ad libitum*. All animals were handled and euthanized according to the animal study protocol approved by the Laboratory Animal Committee, Asahikawa Medical College.

15

### *Media*

All chemicals were obtained from Nacalai Tesque, Inc. (Kyoto, Japan) unless otherwise stated. The medium for oocyte collection and sperm injection was a modified CZB medium (Chatot *et al.*, 1989; Chatot *et al.*, 1990) with 20 mM HEPES, 5 mM  
20 NaHCO<sub>3</sub>, and 0.1 mg/ml polyvinyl alcohol (PVA; cold water soluble; molecular weight: 30,000-70,000, Sigma Chemical Co. Ltd., St. Louis, MO, USA) instead of BSA (termed HEPES-CZB) (Kimura and Yanagimachi, 1995). EGTA medium previously used (Kusakabe *et al.*, 2001) was modified by eliminating NaCl, keeping EGTA constant at 50 mM and increasing the concentration of Tris-HCl. It consisted of 50 mM EGTA and

100 mM Tris-HCl. This medium, called Tris-buffered EGTA, was prepared by mixing 1 ml of 0.5 M EGTA (pH 8.0, adjusted with NaOH) and 1 ml of 1 M Tris-HCl (Trizma®-HCl, pH 7.4, Sigma-Aldrich, Co., St. Louis, MO, USA) and diluting the mixture with water (Water for Embryo Transfer, Sigma-Aldrich, Inc., St. Louis, MO, USA) up to 10 ml. Its final pH was 7.7-7.9. This medium was used within 30 days after preparation.

#### *Collection of mouse spermatozoa*

Dense masses of spermatozoa were collected from the cauda epididymis and placed at the bottom of a 1.5 ml polypropylene microcentrifuge tube (SARSTEDT, Aktiengesellschaft & Co., Numbrecht, Germany) containing 1.2 ml of Tris-buffered EGTA that had been warmed to 37°C. The tube was left standing for 10 min at 37°C to allow spermatozoa swim into the solution. The upper 1 ml of the sperm suspension in the tube was transferred into another tube and was kept in a refrigerator (4°C) for up to 16 days before freeze-drying. Live/dead staining of spermatozoa was performed using FertiLight (Molecular Probes, Inc., Eugene, OR, USA).

#### *Collection of human spermatozoa*

Semen sample collected from each of two healthy men (donors A and B) was allowed to liquefy at 37°C for 30 min. A 0.5 ml aliquot was gently placed at the bottom of a small test tube containing 2 ml of Tris-buffered EGTA that had been pre-warmed to 37°C. The tube was left standing for 10 min at 37°C to allow spermatozoa disperse into the medium. One ml of the upper layer containing the spermatozoa was transferred into another test tube.

### *Freeze-drying*

One-hundred  $\mu$ l of the sperm suspension as prepared above was each put in a 2 ml glass ampoule (Wheaton Scientific, Millville, NJ, USA). Each ampoule was plunged into liquid nitrogen and kept there for 1 min before connected to a lyophilizer (type: FZ2.5, Labconco, Kansas City, MO, USA). After vacuuming for 4 h, each ampoule was flame-sealed and preserved around 4°C (range of temperature: 3°C to 7°C) in a refrigerator. The inside pressure of ampoules at the time of sealing was  $22\text{--}42 \times 10^{-3}$  mbar.

### *Preparation of oocytes*

Females were each injected with 10 IU of pregnant mare's serum gonadotrophin (Teikokuzohki, Tokyo, Japan). Forty-eight hours later, they were injected with 10 IU of human chorionic gonadotrophin (hCG, Mochida, Tokyo, Japan). Mature metaphase II oocytes were collected from oviducts between 15 h and 17 h after hCG injection. They were freed from cumulus cells by treatment with 0.1% bovine testicular hyaluronidase (1000 units/mg, Sigma-Aldrich Co., St. Louis, MO, USA) in the HEPES-CZB medium, then rinsed and kept in modified CZB medium at 37°C for less than 60 min before sperm injection.

### *Intracytoplasmic sperm injection (ICSI)*

ICSI was carried out according to Kimura and Yanagimachi (1995) except that it was performed at 18–25°C. Freeze-dried spermatozoa in each ampoule were rehydrated by adding 100  $\mu$ l of water (Water for Embryo Transfer, Sigma-Aldrich, Inc., St. Louis, MO, USA) immediately after opening the ampoule. All rehydrated spermatozoa were

immotile. They often had broken or missing tails. Those without tails (heads only) were individually injected into each oocyte. Those with broken or “intact” tails were freed from tails by applying a single or a few Piezo pulses to the neck region of the spermatozoon and only the head was injected into each oocyte (Kuretake *et al*, 1996).

Freeze-dried and rehydrated human spermatozoa had grossly “intact” tails (Fig. 4a). As human spermatozoon was much smaller than mouse spermatozoon, the entire body of human spermatozoon was injected into each oocyte. Prior to sperm injection, mouse oocytes were enucleated after 10 min treatment with 5  $\mu$ M cytochalasin B (Wakayama *et al.*, 1998b). Enucleation was done to avoid ethical criticism of creation of an embryo with human and mouse genomes. Theoretically, enucleation is unnecessary because human and mouse chromosomes are markedly different from each other in their structures. ICSI was completed within 1 h after rehydration of freeze-dried mouse and human spermatozoa.

#### *Culture and examination of mouse oocytes injected with spermatozoa*

Sperm-injected mouse oocytes were transferred into droplets (50-100  $\mu$ l) of the modified CZB medium supplemented with 5.56 mM glucose and 5 mg/ml BSA (fraction V, Sigma) under a paraffin oil (Merck KGaA, Darmstadt, Germany) in a Petri dish. The dish was maintained under a humidified atmosphere of 5% CO<sub>2</sub> in air. Eggs with the second polar body and two pronuclei 5-6 h after ICSI were considered normally activated. Eggs at the 2-cell and blastocyst stages were examined 24 and 96 h after ICSI. When an enucleated mouse oocyte was injected with a human spermatozoon, only one large pronucleus appeared in the oocyte. This was the pronucleus of sperm origin. Thus, an oocyte with one pronucleus and one first polar body was considered



activated.

### *Examination of post-implantation development of mouse embryos*

Two-cell mouse embryos were transferred into oviducts of pseudo-pregnant females  
5 that had mated with vasectomized males during the previous night. Because  
chromosome damage induced in male germ cells contributes to early post-implantation  
death (Singer *et al.*, 2006), the number of normally developing fetuses and implantation  
sites were determined in most cases on day 14 of gestation. In practice this provides  
information on the extent of early embryonic loss after implantation, and normal fetuses  
10 at day 14 of pregnancy very rarely fail to develop to full term (Kusakabe *et al.*, 2001).

### *Chromosome analysis*

Between 5 and 6 h after ICSI, sperm-injected oocytes were transferred to a modified  
CZB medium containing 0.01 µg/ml vinblastine sulfate to arrest them at the metaphases  
15 of the first cleavage. Between 20 and 22 h after ICSI, the oocytes were freed from the  
zonae pellucidae by treatment with 0.5% protease (Actinase E, 1000 tyrosine unit/mg,  
Kaken Pharmaceuticals, Tokyo, Japan) followed by 4-10 min treatment with a  
hypotonic solution composed of a 1:1 mixture of 30% fetal bovine serum and 1%  
sodium citrate (Kishikawa *et al.*, 1999; Tateno *et al.*, 2000). The oocytes were fixed by  
20 the gradual fixation method (Kamiguchi and Mikamo, 1986), air-dried on glass slides,  
and stained with Giemsa's solution (Merck KGaA, Darmstadt, Germany) for  
chromosome examination. When mouse chromosomes were examined, the structural  
chromosome aberrations were recorded without discriminating between paternal and  
maternal origins. Because mouse oocytes seldom had chromosome aberrations at the

first mitotic metaphase after normal fertilization and parthenogenetic activation (Tateno and Kamiguchi, 2002), the chromosome groups with aberrant constitution were mostly likely those of sperm origin. Thus, chromosome integrity of mouse spermatozoa could be expressed by the frequency (%) of zygotes with normal chromosome composition (40 acrocentric or telocentric chromosomes) without structural aberrations.

Chromosome integrity of human spermatozoa was expressed by the frequency (%) of the spermatozoa with normal chromosome composition (23 chromosomes) without any structural chromosome aberrations. Of all Giemsa-stained chromosome spreads with normal chromosome composition (semen donor A), twenty were randomly selected and further examined after Q-band staining (Caspersson *et al.*, 1970).

#### *Statistical analysis*

Comparisons of chromosomal integrity of spermatozoa, *in vitro* development and post-implantation development of embryos were made by chi-square analysis using Yate's correction for continuity. Significant differences were accepted with  $P < 0.05$ .

### **Results**

#### *Ability of mouse spermatozoa to activate oocytes after freeze-drying in Tris-buffered EGTA*

The majority of mouse spermatozoa in Tris-buffered EGTA became immotile by 10 min, but they were "alive" (had intact plasma membranes) according to the Live/Dead staining (Fig. 1a). No spermatozoa were alive when they were kept in Tris-buffered EGTA at 4°C for 6 days (Fig. 1b). The spermatozoa freeze-dried after 4-6 days storage

in Tris-buffered EGTA (Fig. 1c) activated oocytes very well, but those kept in the Tris-buffered EGTA for more than 7 days had inferior ability to activate oocytes (Fig. 1d).

#### *Chromosomes of freeze-dried mouse spermatozoa*

Figure 2 shows two typical examples of chromosomes seen on metaphase plates of the first mitotic division (cleavage). Either two groups (a) or one group (b) had normal haploid set (20) of chromosomes. As oocyte chromosomes seldom show abnormalities during normal fertilization and parthenogenetic activation (Tateno and Kamiguchi, 2002), the chromosome group with abnormal constitutions is most likely of sperm origin.

Analysis of chromosomes that appeared on the metaphase plates of the first mitotic division after injection of freeze-dried mouse spermatozoa revealed that the incidence of zygotes with normal chromosomes increased (from 41 to 95%) as the spermatozoa were kept in Tris-buffered EGTA for more than two days before freeze-drying (Table 1). In other words, spermatozoa became more resistant to freeze-drying as they were maintained in Tris-buffered EGTA longer. Chromosome aberrations most commonly found after brief incubation of spermatozoa in Tris-buffered EGTA were chromosome-type breaks or exchanges (Fig. 2b).

#### *Development of oocytes injected with freeze-dried mouse spermatozoa*

Development of mouse oocytes injected with freeze-dried and fresh (non-frozen) spermatozoa was compared. Freeze-dried spermatozoa used in this study were those kept in Tris-buffered EGTA solution for 5-7 days before freeze-drying; the majority of

these spermatozoa had normal chromosomes (Table I). Regardless of the period of storage at 4°C, freeze-dried spermatozoa were as good as fresh (non-frozen) spermatozoa in supporting preimplantation development and even better than fresh (non-frozen) spermatozoa in supporting post-implantation development to mid-term fetuses (Table II, Fig. 3a, b).

#### *Chromosome of freeze-dried human spermatozoa*

When freeze-dried human spermatozoa were rehydrated, their gross morphology was normal (Fig. 4a). Many spermatozoa freeze-dried immediately after the sperm collection and those freeze-dried after pre-freeze-drying incubation had normal chromosomes (Fig. 4b, Table III). Most commonly observed chromosome aberrations were breaks (Table III). Q-Banded 20 chromosome spreads of freeze-dried human spermatozoa did not show any structural chromosome aberrations (Fig. 4c, d).

## **Discussion**

We previously reported that the incidence of mouse spermatozoa with normal chromosomes was lower when they were freeze-dried (68%) than when were frozen-thawed without cryoprotection (89%) (Ward *et al.*, 2003). This was confirmed in the present study. The majority (87-90%) of fresh (non-frozen) or frozen-thawed spermatozoa had normal chromosomes, whereas less than half (41%) of spermatozoa had normal chromosomes when they were freeze-dried after brief suspension in Tris-buffered EGTA (Table 1). Interestingly, the proportion of freeze-dried spermatozoa with normal chromosomes increased (77-95%) when spermatozoa were kept in

Tris-buffered EGTA for more than 2 days before freeze-drying (Table 1). This was quite unexpected because we previously thought that nuclei (chromosomes) of spermatozoa and spermatids became degraded quickly in artificial media when plasma membranes were disrupted or removed (Suzuki *et al.*, 1998; Yanagimachi, 2005). We did not expect  
5 that spermatozoa retained their plasma membranes intact for many days in such an unphysiological solution (Fig. 1b). The reason why prolonged immersion of spermatozoa in Tris-buffered EGTA makes sperm nuclei more resistant to freeze-drying is not clear. EGTA may slowly penetrate into sperm nuclei and chelate intra-nuclear  $\text{Ca}^{2+}$ . During rehydration of freeze-dried spermatozoa, sperm nuclear topoisomerase that  
10 cleaves DNA in the presence of  $\text{Ca}^{2+}$  (Shaman *et al.*, 2006) may be silenced by EGTA. Why do human spermatozoa withstand freeze-drying without long sojourn in Tris-buffered EGTA? It is known that human spermatozoa are unusual in that their nuclear chromatin is less intensely crosslinked by disulphide (S-S) bonds than that of spermatozoa of most other mammals (Bedford *et al.*, 1973). SS-poor human sperm  
15 nuclei may be more readily penetrable by EGTA than SS-rich sperm nuclei of spermatozoa of other species. EGTA we used in this study does chelate  $\text{Ca}^{2+}$ , but not  $\text{Zn}^{2+}$ . Since human sperm chromatin contains 8 mM  $\text{Zn}^{2+}$  (Bjorndahl and Kvist, 2003) and EGTA does not chelate  $\text{Zn}^{2+}$ , EGTA may protect chromatin SH- from oxidizing into superfluous S-S bridges during handling *in vitro* which otherwise could affect sperm  
20 chromatin decondensation and further development into metaphase chromosomes.

In the present study, we examined sperm chromosomes by allowing sperm nuclei transform into metaphase chromosomes of the first mitosis (cleavage) after injection into oocytes. About 10% of non-frozen spermatozoa from normal, fertile mouse males showed chromosome aberrations (Table 1), which was at about the same as previously

reported (Tateno *et al.*, 2000, Ward *et al.*, 2003; Tateno and Kamiguchi 2004). Similarly, about 10% of non-frozen human spermatozoa from normal men showed chromosome aberrations after injection into mouse oocytes (Table 3). This level of sperm chromosome aberration was reported previously after injection of human spermatozoa into mouse oocytes (Lee *et al.*, 1996; Rybouchkin *et al.*, 1996; Watanabe, 2003) or after *in vitro* fertilization of zona-free hamster oocytes with human spermatozoa (Rudak *et al.*, 1978; Martin *et al.*, 1982, 1983; Brandriff *et al.*, 1985; Kamiguchi and Mikamo, 1986; Kusakabe *et al.*, 1996). The real level of chromosome aberrations in mouse and human spermatozoa may be higher or lower depending on individuals. Aberrations at 10% level are likely inherent to the techniques employed (mechanical injection of sperm into oocytes or fusion of sperm with eggs of foreign species). Therefore, more than 80% normality of sperm chromosomes after freeze-drying of mouse and human spermatozoa (Tables 1 and 3) indicates that freeze-drying itself is not detrimental to sperm chromosomes as far as it is properly processed.

Ability of mouse spermatozoa to activate oocytes decreased after the pre-freeze-drying incubation for more than 7 days in Tris-buffered EGTA (Fig. 1d). When mouse spermatozoa were kept in Tris-buffered EGTA solution for more than 6 days, sperm's ability to activate oocytes decreased (Fig. 1d). This is likely due to deterioration of sperm-born oocyte-acting factors (SOAF). Nature of mammalian SOAF has been the subject of intense studies. Thus far, strong candidates include phospholipase C zeta (Cox *et al.*, 2002; Saunders *et al.*, 2002), truncated c-kit tyrosine kinase (Sette *et al.*, 1997, 2002) and WW domain-binding protein (Wu *et al.*, 2007). At least part of it is localized in the perinuclear theca (Kimura *et al.*, 1998; Wu *et al.*, 2007). SOAF may not be very stable in EGTA-buffer, but once it is freeze-dried, it seems to

retain its biological activity for long time.

Freeze-dried mouse spermatozoa may maintain their chromosome integrity and developmental potential indefinitely at -80°C (Kawase *et al.*, 2005) or at -196°C (liquid nitrogen), but constant need of deep freezers or liquid nitrogen is troublesome.

5 According to Kaneko and Nakagata (2006), mouse spermatozoa freeze-dried in EDTA (ethylenediaminetetraacetic acid) solution (1 mM EDTA + 10 mM Tris-HCl) well maintained chromosome integrity. Freeze-dried spermatozoa which were kept at 4°C for 1 year were able to produce live offspring at the about the same rate as we reported in the present paper. These investigators did not study whether sperm incubation in EDTA  
10 solution renders sperm chromosomes more resistant to freeze-drying. Our ultimate goal is to maintain freeze-dried mammalian spermatozoa at ambient temperature without using refrigerator/freezer or dry ice/liquid nitrogen.

### Acknowledgement

15 This work was supported by Grants-in-Aid for Scientific Research, No. 16681020 from the Ministry of Education, Culture, Sports, Science and Technology of Japan (to H.K.) and by a grant from The Akiyama Foundation (to H.K.).

## References

- Bedford JM, Calvin H and Cooper GW (1973) The maturation of spermatozoa in the human epididymis. *J Reprod Fertil Suppl* 18,199-213.
- Bhowmick S, Zhu L, McGinnis L, Lawitts J, Nath BD, Toner M and Biggers J (2003) Desiccation tolerance of spermatozoa dried at ambient temperature: production of fetal mice. *Biol Reprod* 68,1779-1786.
- Brandriff B, Gordon L, Ashworth L, Watchmaker G, Moore D 2nd, Wyrobek AJ and Carrano AV (1985) Chromosomes of human sperm: variability among normal individuals. *Hum Genet* 70,18-24.
- 10 Bjorndahl L and Kvist U (2003) Sequence of ejaculation affects the spermatozoon as a carrier and its message. *Reprod Biomed Online* 7,440-448.
- Caspersson T, Zech L and Johansson C (1970) Differential binding of alkylating fluorochromes in human chromosomes. *Exp Cell Res* 60,315-319.
- Chatot CL, Ziomek A, Bavister BD, Lewis JL and Torres I (1989) An improved culture medium supports development of random-bred 1-cell mouse embryos in vitro. *J Reprod Fertil* 86,679-688.
- 15 Chatot CL, Lewis JL, Torres I and Ziomek CA (1990) Development of 1-cell embryos from different strains of mice in CZB medium. *Biol Reprod* 42,432-440.
- Cox LJ, Larman MG, Saunders CM, Hashimoto K, Swann K and Lai FA (2002) Sperm phospholipase C $\zeta$  from humans and cynomolgus monkeys triggers Ca<sup>2+</sup> oscillations, activation and development of mouse oocytes. *Reproduction* 124,611-623.
- 20 Kamiguchi Y and Mikamo K (1986) An improved, efficient method for analyzing human sperm chromosomes using zona-free hamster ova. *Am J Hum Genet*



38,724-740.

Kaneko T and Nakagata N (2006) Improvement in the long-term stability of freeze-dried mouse spermatozoa by adding of a chelating agent. *Cryobiology* 53,279-282.

- 5 Kawase Y, Araya H, Kamada N, Jishage K and Suzuki H (2005) Possibility of long-term preservation of freeze-dried mouse spermatozoa. *Biol Reprod* 72,568-573.

Kimura Y and Yanagimachi R (1995) Intracytoplasmic sperm injection in the mouse. *Biol Reprod* 52,709-720.

- Kimura Y, Yanagimachi R, Kuretake S, Bortkiewicz H, Perry AC and Yanagimachi H  
10 (1998) Analysis of mouse oocyte activation suggests the involvement of sperm perinuclear material. *Biol Reprod* 58,1407-1415.

Kishikawa H, Tateno H and Yanagimachi R (1999) Chromosome analysis of BALB/c mouse spermatozoa with normal and abnormal head morphology. *Biol Reprod* 61,809-812.

- 15 Kuretake S, Kimura Y, Hoshi K and Yanagimachi R (1996) Fertilization and development of mouse oocytes injected with isolated sperm heads. *Biol Reprod* 55,789-795.

- Kusakabe H, Yamakage K and Tanaka N (1996) Detection of neocarzinostatin-induced translocations in human sperm chromosomes using  
20 fluorescence in situ hybridization of chromosome 2. *Mutat Res* 369,51-58.

Kusakabe H, Szczygiel MA, Whittingham DG and Yanagimachi R (2001) Maintenance of genetic integrity in frozen and freeze-dried mouse spermatozoa. *Proc Natl Acad Sci USA* 98,13501-13506.

Lee JD, Kamiguchi Y and Yanagimachi R (1996) Analysis of chromosome constitution

of human spermatozoa with normal and aberrant head morphologies after injection into mouse oocytes. Hum Reprod 11,1942-1946.

Li MW, Biggers JD, Elmoazzen HY, Toner M, McGinnis L and Lloyd KC (2007)

Long-term storage of mouse spermatozoa after evaporative drying. Reproduction  
5 (in press).

Martin RH, Lin CC, Balkan W and Burns K (1982) Direct chromosomal analysis of human spermatozoa: preliminary results from 18 normal men. Am J Hum Genet 34,459-468.

Martin RH, Balkan W, Burns K, Rademaker AW, Lin CC and Rudd NL (1983) The  
10 chromosome constitution of 1000 human spermatozoa. Hum Genet 63,305-309.

McGinnis LK, Zhu L, Lawitts JA, Bhowmick S, Toner M and Biggers JD (2005) Mouse sperm desiccated and stored in trehalose medium without freezing. Biol Reprod 73,627-633.

Nakagata N (2000) Cryopreservation of mouse spermatozoa. Mamm Genome  
15 11,572-576.

Ogonuki N, Mochida K, Miki H, Inoue K, Fray M, Iwaki T, Moriwaki K, Obata Y, Morozumi K, Yanagimachi R *et al.* (2006) Spermatozoa and spermatids retrieved from frozen reproductive organs or frozen whole bodies of male mice can produce normal offspring. Proc Natl Acad Sci USA 103,13098-13103.

20 Polge C, Smith AU and Parks AS (1949) Revival of spermatozoa after vitrification and dehydration at low temperatures. Nature 164,666.

Rudak E, Jacobs PA and Yanagimachi R (1978) Direct analysis of the chromosome constitution of human spermatozoa. Nature 274,911-913.

Rybouchkin A, Dozortsev D, Pelinck MJ, De Sutter P and Dhont M. (1996) Analysis of

the oocyte activating capacity and chromosomal complement of round-headed human spermatozoa by their injection into mouse oocytes. Hum Reprod 11,2170-2175.

Saunders CM, Larman MG, Parrington J, Cox LJ, Royse J, Blayney LM, Swann K and

- 5 Lai FA (2002) PLC zeta: a sperm-specific trigger of  $Ca^{2+}$  oscillations in eggs and embryo development. Development 129,3533-3544.

Sette C, Bevilacqua A, Bianchini A, Mangia F, Geremia R and Rossi P (1997)

Parthenogenetic activation of mouse eggs by microinjection of a truncated c-kit tyrosine kinase present in spermatozoa. Development 124,2267-2274.

- 10 Sette C, Paronetto MP, Barchi M, Bevilacqua A, Geremia R and Rossi P. (2002)

Tr-kit-induced resumption of the cell cycle in mouse eggs requires activation of a Src-like kinase. EMBO J 21,5386-5395.

Shaman JA, Prisztoka R and Ward WS (2006) Topoisomerase IIB and an extracellular nuclease interact to digest sperm DNA in an apoptotic-like manner. Biol Reprod

- 15 75,741-748.

Singer TM, Lambert IB, Williams A, Douglas GR and Yauk CL (2006) Detection of induced male germline mutation: correlations and comparisons between traditional germline mutation assays, transgenic rodent assays and expanded simple tandem repeat instability assays. Mutat Res 598,164-193.

- 20 Suzuki K, Yanagida K and Yanagimachi R (1998) Comparison of the media for isolation and storage of round spermatid nuclei before intracytoplasmic injection. J Assist Reprod Genet 15,154-157.

Tateno H, Kimura Y and Yanagimachi R (2000) Sonication per se is not as deleterious to sperm chromosomes as previously inferred. Biol Reprod 63,341-346.

Tateno H and Kamiguchi Y (2002) Abnormal chromosome migration and chromosome aberrations in mouse oocytes during meiosis II in the presence of topoisomerase II inhibitor ICRF-193. *Mutat Res* 502,1-9.

Tateno H and Kamiguchi Y (2004) Chromosome analysis of mouse one-cell

5 androgenones derived from a sperm nucleus exposed to topoisomerase II inhibitors at pre- and post-fertilization stages. *Mutat Res* 556,117-126.

Wakayama T, Whittingham DG and Yanagimachi R (1998a) Production of normal offspring from mouse oocytes injected with spermatozoa cryopreserved with or without cryoprotection. *J Reprod Fertil* 112,11-17.

10 Wakayama T, Perry ACF, Zuccotti M, KR Johnson KR and Yanagimachi R (1998b) Full-term development of mice from enucleated oocytes injected with cumulus cell nuclei. *Nature* 394,369-374.

Wakayama T and Yanagimachi R (1998) Development of normal mice from oocytes injected with freeze-dried spermatozoa. *Nat Biotechnol* 16,639-641.

15 Ward MA, Kaneko T, Kusakabe H, Biggers JD, Whittingham DG and Yanagimachi R (2003) Long-term preservation of mouse spermatozoa after freeze-drying and freezing without cryoprotection. *Biol Reprod* 69,2100-2108.

Watanabe S (2003) A detailed cytogenetic analysis of large numbers of fresh and frozen-thawed human sperm after ICSI into mouse oocytes. *Hum Reprod*  
20 18,1150-1157.

Wu AT, Sutovsky P, Manandhar G, Xu W, Katayama M, Day BN, Park KW, Yi YJ, Xi YW, Prather RS *et al.* (2007) PAWP, a sperm-specific WW domain-binding protein, promotes meiotic resumption and pronuclear development during fertilization. *J Biol Chem* 282,12164-12175.

Yanagimachi R (2005) Intracytoplasmic injection of spermatozoa and spermatogenic cells: its biology and applications in humans and animals. Reprod Biomed Online 10,247-288.

5

10

15

20

## Legend of Figures

Figure 1. Freeze-dried mouse spermatozoa after Live/Dead staining. (a) Most spermatozoa immediately after suspension in Tris-buffered EGTA were stained green (alive), (b) whereas those kept in the same solution for 6 days at 4°C were stained red (dead). (c) Freeze-dried spermatozoa after rehydration; most had broken tails; heads of some spermatozoa were completely separated from tails. (d) A diagram showing that the ability of freeze-dried mouse spermatozoa to activate oocytes by ICSI decreases when pre-freeze-drying incubation of spermatozoa prior to freeze-drying becomes longer than 7 days. The values are expressed as mean  $\pm$  SD.

Figure 2. Chromosome spreads of oocytes injected with freeze-dried mouse spermatozoa. (a) Chromosomes from two pronuclei (of sperm and oocyte origin) have normal chromosomes. (b) A group of chromosomes with abnormal constitution is most likely of sperm origin. Small arrows indicate chromosome fragments derived from chromosome break or chromosome exchange. A large arrow indicates chromosome exchange.

Figure 3. Embryos developed from oocytes injected with freeze-dried mouse spermatozoa. (a) Blastocysts developed from the oocytes injected with spermatozoa freeze-dried after incubation in Tris-buffered EGTA for 6 days. Arrows indicate the degenerated embryos. (b) Seven normal fetuses and six implantation sites at day 14 of gestation, developed from thirteen 2-cell embryos transplanted into a surrogate mother. These 2-cell embryos were developed from oocytes injected with mouse spermatozoa

freeze-dried after incubation in Tris-buffered EGTA for 6 days.

Figure 4. Freeze-dried human spermatozoa and their chromosomes. (a) Freeze-dried spermatozoa rehydrated. (b) Human sperm chromosomes with normal constitution that  
5 appeared at the first mitotic metaphase after injection of a single freeze-dried spermatozoon into an enucleated mouse oocyte. (c) and (d), Normal Q-banded chromosomes of a freeze-dried X-bearing spermatozoon (c) and a Y-bearing spermatozoon (d).

Table I. Chromosome integrity of mouse zygotes produced by injection of oocytes with frozen and freeze-dried spermatozoa

Mouse spermatozoa injected	Pre-freeze-drying incubation in Tris-buffered EGTA soln. (4°C)	Storage period of freeze-dried spermatozoa (4°C)	No. zygotes analyzed (Exp. No.)	No. (%) zygotes with normal chromosome composition [range]
Fresh (non-frozen) <sup>a</sup>	-	-	80 (3)	72 ( 90 ) <sup>d</sup> [84-92]
Frozen-thawed <sup>b</sup>	-	-	75 (3)	65 ( 87 ) <sup>d, e</sup> [81-100]
Freeze-dried <sup>c</sup>	0 min	1 - 3 d	78 (4)	32 ( 41 ) <sup>f</sup> [36-53]
Freeze-dried	1 d	1 - 3 d	75 (3)	33 ( 44 ) <sup>f</sup> [35-50]
Freeze-dried	2 d	2 - 5 d	82 (3)	63 ( 77 ) <sup>e</sup> [75-78]
Freeze-dried	3 - 4 d	4 - 5 d	121 (6)	107 ( 88 ) <sup>d</sup> [79-95]
Freeze-dried	5 - 7 d	2 - 14 d	93 (4)	79 ( 85 ) <sup>d</sup> [81-89]
Freeze-dried	5 - 7 d	2 - 4 mo	93 (5)	88 ( 95 ) <sup>d</sup> [85-100]

<sup>a</sup> Spermatozoa were suspended in (allowed to swim into) Tris-buffered EGTA solution for 10 min at 37°C before injection into oocytes.

<sup>b</sup> Spermatozoa were suspended in Tris-buffered EGTA solution for 10 min at 37°C was snap-frozen with liquid nitrogen, then thawed before injection into oocytes.

<sup>c</sup> Spermatozoa were suspended in Tris-buffered EGTA solution for 10 min at 37°C before freeze-drying.

<sup>d-f</sup> Significantly different values ( $P$ , 0.05) between different superscripts



Table II. Development of mouse oocytes injected with spermatozoa which were freeze-dried after 5-7 days of incubation in Tris-buffered EGTA (4°C)

Mouse spermatozoa injected	Storage period of freeze-dried spermatozoa (4°C)	No. oocytes survived and activated after sperm injection (No. exp.)	No. (%) oocytes developed into:		No. 2-cell embryos transferred (No. recipients)	Examination on day 14 of gestation	
			2-Cell embryos	Blastocysts		No. (%) implants <sup>a</sup>	No. (%) normal fetuses [range, %]
Fresh (non-frozen) <sup>b</sup>	-	111 ( 5 )	111 ( 100 ) <sup>c</sup>	90 ( 81 ) <sup>c</sup>	-	-	-
Fresh (non-frozen) <sup>b</sup>	-	85 ( 4 )	82 ( 96 ) <sup>c,d</sup>	-	76 ( 5 )	56 ( 74 ) <sup>c</sup>	33 ( 43 ) <sup>c</sup> [13-69]
Freeze-dried	2 - 6 wk	119 ( 4 )	116 ( 97 ) <sup>c,d</sup>	97 ( 82 ) <sup>c</sup>	-	-	-
Freeze-dried	4 d - 3 wk	66 ( 3 )	64 ( 97 ) <sup>c,d</sup>	-	63 ( 4 )	56 ( 89 ) <sup>d</sup>	38 ( 60 ) <sup>c,d</sup> [44-77]
Freeze-dried	3 mo	101 ( 4 )	97 ( 96 ) <sup>c,d</sup>	85 ( 84 ) <sup>c</sup>	-	-	-
Freeze-dried	3 mo	76 ( 4 )	71 ( 93 ) <sup>d</sup>	-	71 ( 4 )	62 ( 87 ) <sup>c,d</sup>	41 ( 58 ) <sup>c,d</sup> [40-76]
Freeze-dried	1 yr	88 ( 3 )	84 ( 95 ) <sup>c,d</sup>	-	84 ( 6 )	74 ( 88 ) <sup>d</sup>	54 ( 64 ) <sup>d</sup> [50-77] <sup>e</sup>

<sup>a</sup> Absorbed fetuses plus normal fetuses

<sup>b</sup> Spermatozoa were suspended in (allowed to swim into) Tris-buffered EGTA solution for 10 min at 37°C.

<sup>c,d</sup> Significantly different values (*P* , 0.05) between different superscripts within each column

<sup>e</sup> Examined on day 18 of gestation.

Table III. Chromosome analysis of freeze-dried human spermatozoa after injection into enucleated mouse oocytes

Semen donor	Pre-freeze-drying sperm incubation in Tris-buffered EGTA (4°C)	Storage period of freeze-dried spermatozoa (4°C)	No. of oocytes injected with spermatozoa (No. exp)	No. (%) oocytes activated and reached metaphase of first mitosis	No. (%) oocytes with normal sperm chromosome	No. spermatozoa with chromosome aberrations						
						Structural aberrations					Aneuploidy	
						ctb	cte	csb	cse	mul	hyper	hypo
A	0 min <sup>a</sup>	1 d	71 (4)	63 ( 89 ) <sup>b</sup>	56 ( 89 ) <sup>d</sup>	0	0	7	0	0	0	0
A	0 min <sup>a</sup>	3 - 7 d	71 (5)	61 ( 86 ) <sup>b, c</sup>	58 ( 95 ) <sup>d</sup>	0	0	3	0	0	0	0
A	0 min <sup>a</sup>	4 wk	74 (5)	67 ( 91 ) <sup>b</sup>	63 ( 94 ) <sup>d</sup>	1	0	2	1	0	0	0
A	6 - 7 d	3 wk	78 (5)	57 ( 73 ) <sup>c</sup>	50 ( 88 ) <sup>d</sup>	0	0	6	1	0	0	0
B	0 min <sup>a</sup>	3 d - 4 wk	62 (4)	57 ( 92 ) <sup>b</sup>	49 ( 86 ) <sup>d</sup>	0	0	6	1	1	0	0
B	4 h	4 d - 4 wk	66 (4)	58 ( 88 ) <sup>b, c</sup>	49 ( 84 ) <sup>d</sup>	1	2	5	1	0	1	0
B	2 d	17 d - 4 wk	56 (4)	41 ( 73 ) <sup>c</sup>	33 ( 80 ) <sup>d</sup>	0	1	5	0	2	0	0

<sup>a</sup> Spermatozoa were suspended in (allowed to swim into) Tris-buffered EGTA solution for 10 min at 37°C before freeze-drying.

<sup>b, c</sup> Significantly different values ( $P$ , 0.05) between different superscripts within the data from "A" or "B"

<sup>d</sup> Not significant difference ( $P$ , 0.05) within the data from "A" or "B"

Abbreviations, ctb: chromatid break; cte: chromatid exchange; csb: chromosome break; cse: chromosome exchange; mul: multiple aberrations; hyper: hyperploidy; hypo: hypoploidy.

Figure 1





Figure 3

a



b




Figure 2




Figure 4

