

# AMCoR

Asahikawa Medical University Repository <http://amcor.asahikawa-med.ac.jp/>

Biological and Pharmaceutical Bulletin (2014) 37(12):1907–1912.

Co-localization of TRPV2 and Insulin-Like Growth Factor-I Receptor in  
Olfactory Neurons in Adult and Fetal Mouse

Hitoshi Matsui, Tomohiro Noguchi, Kaoru Takakusaki,  
Makoto Kashiwayanagi

## Co-localization of TRPV2 and Insulin-Like Growth Factor-I Receptor in Olfactory Neurons in Adult and Fetal Mouse

Hitoshi Matsui, Tomohiro Noguchi, Kaoru Takakusaki, and Makoto Kashiwayanagi\*

Department of Sensory Physiology, Asahikawa Medical University, Midorigaoka E2-1, Asahikawa 078-8510, Japan.

Received June 3, 2014; accepted September 7, 2014

TRPV2, a member of the transient receptor potential family, has been isolated as a capsaicin-receptor homolog and is thought to respond to noxious heat. Here we show that TRPV2 mRNA is predominantly expressed in the subpopulation of olfactory sensory neurons (OSNs). We carried out histochemical analyses of TRPV2 and insulin-like growth factor-I receptor (IGF-IR) using *in situ* hybridization and immunofluorescence in the adult olfactory system. In olfactory mucosa, intensive TRPV2 immunostaining was observed at the olfactory axon bundles but not at the soma. TRPV2-positive labeling was preferentially found in the olfactory nerve layer in the olfactory bulb (OB). Furthermore, we demonstrated that a positive signal for IGF-IR mRNA was detected in OSNs expressing TRPV2 mRNA. In embryonic stages, TRPV2 immunoreactivity was observed on axon bundles of developing OSNs in the nasal region starting from 12.5 d of gestation and through fetal development. Observations in this study suggest that TRPV2 coupled with IGF-IR localizes to growing olfactory axons in the OSNs.

**Key words** olfactory neuron; olfactory epithelium; olfactory bulb; insulin-like growth factor-I receptor; transient receptor potential V2

Olfactory sensory neurons (OSNs), which receive odorants, transmit odor information to the olfactory bulb (OB). Some volatile odorants having toxic effects physiologically damage the OSNs. In addition, because the olfactory organ is located at the peripheral region, physical damage is likely to occur in the OSNs. One of their particular features is that they are continuously renewed during adulthood to overcome these disadvantages.<sup>1,2</sup> The life period of rat OSNs was about 30 d. To transmit odor information to the brain, newly generated OSNs extend their axons to the OB to form synaptic connections with mitral/tufted cells at the glomeruli of the OB throughout an individual's life.

It has been suggested that the proliferation and differentiation of OSNs are controlled by several trophic factors.<sup>3–9</sup> In the rodent or avian olfactory system, insulin-like growth factor (IGF)-I is produced in the OB at the embryonic and postnatal stages,<sup>10,11</sup> while the IGF-I receptor (IGF-IR) is expressed in both OB and olfactory epithelium (OE).<sup>12</sup> These results suggest that IGF-I plays an important role in bulbo-epithelial interactions in the olfactory system.

In general, axon elongation and guidance depend on the localization, concentration, and temporal dynamics of cytoplasmic calcium signaling.<sup>13,14</sup> Whereas increases in intracellular calcium levels lead to the activation of calcium-dependent signaling molecules, the location of a regulatory calcium-permeable channel remains unclear, as does the way in which its activity is regulated. Transient receptor potential (TRP) ion channels form a large superfamily of non-selective cation channels that display a remarkable diversity of activation mechanisms.<sup>15</sup> Members of the vanilloid receptor family (TRPV) are activated by a wide range of stimuli, including temperature, protons, phorbol esters, lipids, changes in extracellular osmolarity and pressure, and depletion of intracellular calcium stores. A trophic factor such as nerve growth factor increases TRPV1, which is activated by an increase in temperature or capsaicin,<sup>16</sup> functions *via* phosphatidylinositol

3-kinase (PI3K)-regulated trafficking to the plasma membrane.<sup>17</sup> TRPV2 (also known as VRL-1 or GRC), a TRPV1 homologue, is activated by high-intensity noxious heat.<sup>16</sup> In a preliminary experiment, reverse-transcription polymerase chain reaction (RT-PCR) analysis using specific primers for each 22 plasma membrane-associated TRP family, TRPC1-7, TRPM1-8, TRPV1-6, and TRPA1 in mouse olfactory mucosa clearly detected amplification of fragment for TRPC1, TRPC6, TRPM3, TRPM4, TRPM5, TRPM6, TRPM7, TRPV2, TRPV6, and TRPA1 (data not shown). *In situ* hybridization showed that cell bodies labeled with molecular probes selectively directed against TRPM3, TRPM4, TRPM5, TRPM6, TRPM7, TRPV2, TRPV6, and TRPA1 mRNAs were located in the olfactory epithelium (data not shown). Ahmed *et al.*<sup>18</sup> and Nakashimo *et al.*<sup>19</sup> support that TRPV2 is expressed in mouse olfactory epithelium.

Kanzaki *et al.*<sup>20</sup> have reported that translocation of TRPV2 to the plasma membrane is augmented by IGF-I, although the physiological significance of IGF-I-regulated TRPV2 sensitization remains unknown. In this study, we carried out histochemical analyses of TRPV2 and IGF-I receptor using *in situ* hybridization and immunofluorescence in the adult olfactory system to explore the essential calcium channel that regulates axon elongation and guidance. We also investigated the immunohistochemical localization of TRPV2 in the OSNs of the nasal regions during fetal development.

### MATERIALS AND METHODS

**Antibodies** Purified rabbit anti-TRPV2 polyclonal immunoglobulin G (IgG), which was raised against amino acids 744–761 (KNSASEEDHLPLQLVQSP) of TRPV2, was obtained from Oncogene Research Products (San Diego, CA, U.S.A.). Mouse monoclonal antibody against  $\beta$ III-tubulin and growth-associated protein-43 (GAP43) were obtained from Promega (Tokyo, Japan) and Sigma-Aldrich (Tokyo, Japan), respectively. Olfactory marker protein (OMP) antiserum immunized in goat was a gift from Dr F. Margolis (University

The authors declare no conflict of interest.

\* To whom correspondence should be addressed. e-mail: yanagi@asahikawa-med.ac.jp

of Maryland School of Medicine).<sup>21)</sup> Alexa Fluor 488 donkey anti-rabbit IgG (H+L) and Alexa Fluor 546 donkey anti-goat IgG (H+L) were purchased from Molecular Probes (Tokyo, Japan). Rhodamine-conjugated donkey anti-mouse IgG (H+L) was from Alexis (Tokyo, Japan)

**Animals and Tissue Sample Collection for *in Situ* Hybridization and Immunohistochemistry** The adult (8–12 weeks old) C57BL/6 male mice used for this study were purchased from Sankyo Laboratory (Sapporo, Japan). The mice were kept in a room at  $22 \pm 0.5^\circ\text{C}$  and 58% relative humidity, with a 12h light : 12h dark cycle (light off at 20:00h). All mice had free access to food and water in the housing cage. The animal-use protocol was approved by the Asahikawa Medical University Animal Experiment Committee, and we followed the Asahikawa Medical University Guidelines for Animal Experiments throughout the study. Tissues were immediately snap-frozen in liquid nitrogen and stored at  $-80^\circ\text{C}$  until use. To prepare the frozen sections for immunofluorescence and *in situ* hybridization, mice were deeply anesthetized by intraperitoneal injection of pentobarbital sodium (50mg/kg), followed by intracardial perfusion with 4% paraformaldehyde in phosphate-buffered saline (PBS). Extracted nasal tissues were decalcified by agitating in 0.5M ethylenediaminetetraacetic acid (EDTA)/PBS at  $4^\circ\text{C}$  for 3d. Timed pregnant mice at embryonic days (E) 10.5, 11.5, 12.5, 14.5, and 17.5 were fixed in 4% paraformaldehyde and stored in PBS until use. Tissues were cryoprotected by incubation in 30% sucrose/PBS at  $4^\circ\text{C}$  for 12h, embedded in O.C.T compound, and cryosectioned ( $12\mu\text{m}$ ) using a JUNG CM3000 or CM3050 cryostat (Leica, Nussloch, Germany).

***In Situ* Hybridization** Antisense and sense probes were created by incubating  $1\mu\text{g}$  of linearized plasmid DNA harboring 346 base-pair mouse TRPV2 cDNA fragments (accession no. NM\_011706, 2434–2779) obtained by polymerase chain reaction (PCR) with a 10×digoxigenin (DIG) RNA labeling mixture (Roche, Tokyo, Japan) and T7 or SP6 RNA polymerase (Roche) for 2h at  $37^\circ\text{C}$ . *In situ* hybridization analysis against olfactory epithelium was performed according to a previously described protocol.<sup>22)</sup> Briefly, cryosections were post-fixed in 4% paraformaldehyde in PBS for 10min, washed twice with PBS, acetylated for 10min with 0.25% acetic anhydride in 100mM triethanolamine-HCl (pH 8.0) and again washed twice with PBS. Prehybridization was performed at room temperature with 200  $\mu\text{L}$  hybridization buffer per slide for 3h. The hybridization buffer was composed of 50% formamide (Wako Pure Chemical Industries, Ltd., Osaka, Japan), 5×saline-sodium citrate (SSC) buffer (20×SSC; 3M NaCl, 0.3M sodium citrate, pH 7.0), 5×Denhardt's (Wako Pure Chemical Industries, Ltd.), 250  $\mu\text{g}/\text{mL}$  yeast tRNA (Roche), and 500  $\mu\text{g}/\text{mL}$  herring sperm DNA (Roche). The hybridization was carried out in hybridization buffer containing 200 ng/mL DIG-cRNA for 16h at  $72^\circ\text{C}$ . Slides were washed three times for 20min each with  $0.2 \times \text{SSC}$  at  $72^\circ\text{C}$ . The hybridized DIG-labeled probe was detected using alkaline phosphatase-conjugated anti-DIG antibody (Roche) at a dilution of 1:5000 and visualized with nitro-blue tetrazolium chloride (NBT)/5-bromo-4-chloro-3'-indolylphosphatase *p*-toluidine salt (BCIP).

**Double-Fluorescent *In Situ* Hybridization** The simultaneous expression of two genes, IGF-IR and TRPV2, was detected using DIG-labeled IGF-IR RNA probe (NM\_010513, 5831–7210bp) and fluorescein-labeled TRPV2 RNA probe

(NM\_011706, 395–1435bp). These probes were created according to above, except for the fluorescein labeling with a 10× fluorescein RNA labeling mix (Roche, Mannheim, Germany). Hybridizing and washing conditions were same as described above, except for applying of both DIG- and fluorescein-labeled probes to  $5\mu\text{m}$  coronal cryosections of olfactory epithelium. For amplification of the signal of the fluorescein-labeled probe, TSA-plus fluorescein system (PerkinElmer, Inc., Waltham, MA, U.S.A.) was employed after blocking with 1% blocking reagent (Roshe) for 1h and incubation with 1/400 anti-fluorescein antibody conjugated HRP (PerkinElmer, Inc.) for 1h at  $37^\circ\text{C}$ . In addition, the tyramide-fluorescein signal was strengthened to Alexa Fluor 488 signal by two incubations, with 1/100 rabbit anti-fluorescein (AbD serotec, Oxford, U.K.) for 1h at  $37^\circ\text{C}$  and with 1/50 donkey anti-rabbit IgG conjugated Alexa Fluor 488 (Molecular Probes, Eugene, OR, U.S.A.) for 1h at  $37^\circ\text{C}$ . Each of the incubations with the antibodies was followed by washing with PBS containing 0.05% tween 20. The DIG-labeled signal detected with 1/5000 alkaline phosphatase-conjugated anti-DIG antibody was visualized with HNPP fluorescent detection set (Roche) according to the manufacturer's instruction.

**Immunofluorescence** Air-dried cryosections or cells plated on a glass-bottom dish were incubated in PBS containing 0.5% Triton X-100 and 1% bovine serum albumin (BSA) (blocking solution), followed by overnight incubation at  $4^\circ\text{C}$  in blocking solution containing primary antibody. The sections were subsequently washed, and a positive immunoreaction was visualized by using Alexa Fluor 488 donkey anti-rabbit IgG, rhodamine-conjugated donkey anti-mouse IgG, and Alexa Fluor 546 donkey anti-goat IgG, diluted to 1:100 in blocking solution for 1h. After washing, sections were counterstained with DAPI (Dojindo, Kumamoto, Japan) and mounted with Aqua Poly/Mount (Polysciences, Warrington, PA, U.S.A.). Microphotographs were taken using fluorescent optics on an IX71 inverted microscope (Olympus, Tokyo, Japan) with a C4742-95 camera (Hamamatsu Photonics, Hamamatsu, Japan).

## RESULTS

### Cellular Localization of TRPV2 mRNA in Adult Mouse Olfactory Mucosa

The olfactory mucosa consists of heterogeneous cells such as olfactory receptor, supporting, and basal cells. To determine the cellular localization of TRPV2 in olfactory mucosa, we characterized the expression patterns of TRPV2 mRNAs in the olfactory mucosa using *in situ* hybridization (Fig. 1). The general cytoarchitecture of olfactory epithelium (OE) with a laminar distribution according to the level of differentiation<sup>1,2)</sup> is shown in the Fig. 1a staining for olfactory marker protein (OMP), a marker for mature olfactory sensory neuron (OSN).<sup>21)</sup> Olfactory stem cells (called basal cells) and mitotic progenitors lie in the deep OE. Several layers of mature OSNs that express OMP are present in the middle compartment of the OE. As shown in Fig. 1b, cell bodies labeled with molecular probes selectively directed against TRPV2 abounded in OSNs of the OE. In the OE, the supporting cells formed a single superficial layer clearly lacking TRPV2 mRNAs. No significant labeling in *in situ* hybridization analysis could be obtained with the sense probes (not shown).

### Immunolocalization of TRPV2 in Adult Mouse Olfac-

**tory Mucosa and Olfactory Bulb** To determine the precise distribution of TRPV2 protein, we performed double-labeling immunofluorescence using antibodies against TRPV2 and OMP or growth-associated protein-43 (GAP43). Just above this level of proliferation, postmitotic immature OSNs are known to express GAP43<sup>23</sup> and are OMP-negative. Olfactory nerve bundles lying in the lamina propria displayed intense anti-TRPV2 fluorescence (Figs. 2a, d). TRPV2-positive bundles were co-localized with OMP (Figs. 2b, c). TRPV2 immunoreactivities were also observed in axon bundles labeled with GAP43 (Figs. 2e, f). However, obvious staining on cell bodies and cilia of the OSNs was not found by immunofluorescence with the antibody against TRPV2. This result suggests that most TRPV2 protein is quickly transported to axons of GAP43-positive immature and OMP-positive mature OSNs, and that protein amounts in soma are too low to be detected by our immunofluorescence.

The axons of the OSNs from the olfactory mucosa form an OMP-positive olfactory nerve layer (ONL in Fig. 3a) at the surface of the olfactory bulb (OB). Consistent with the intense TRPV2 immunolabeling observed in the OMP-positive olfac-

tory nerve bundles lying in the deep OE, the glomerular layer and olfactory nerve layer of the OB displayed immunoreactivity for TRPV2 (Fig. 3b). More abundant axons labeled with TRPV2 antibody were found to be concentrated in the olfactory nerve layer than in glomeruli, suggesting that TRPV2 localizes on olfactory axons and that the TRPV2-positive axons reach their target glomeruli.

**TRPV2 mRNA and Insulin-Like Growth Factor-I Receptor mRNA Expression in Mouse OSNs** Previous reports have indicated that translocation of TRPV2 is augmented by insulin-like growth factor (IGF)-I through PI3K and the Akt signaling pathway, leading to a gradual increase in calcium permeability.<sup>18</sup> To explore the presence of the IGF system in olfactory mucosa, we examined the existence of mRNA of the IGF-IR in the OE (Fig. 4). As shown in Fig. 1, subpopulation of OSNs expressed TRPV2 mRNA (Fig. 4a, green). Cell bodies of labeled with molecular probes selectively directed against IGF-I receptor abundant in OSNs (Fig. 4b, red). Most OSNs expressing TRPV2 mRNA expressed also IGF-IR mRNA (Fig. 4c, yellow).

**TRPV2 Immunolocalization during Fetal Development** If TRPV2 is confined to elongating axons of newly differentiated OSN, as expected from the localization of TRPV2 in the adult OB, TRPV2 may regulate the extension of axons of the OSNs in nasal regions during fetal development. To determine whether or not TRPV2 is involved in olfactory system development, we examined immunofluorescence using E10.5, E11.5, E12.5, E14.5, and E17.5 embryos. No significant signals in the nasal region were observed by immunofluorescence using antibodies against TRPV2 at E10.5 and E11.5 (not shown). We detected TRPV2-immunopositive staining in axon bundles labeled with  $\beta$ III-tubulin, a marker of newly differentiated neurons<sup>24</sup> at E12.5 (Figs. 5a–c), when OMP is still not expressed in the OSNs. This observation indicated that TRPV2 localizes to axons of immature OSNs in the fetal development. At E14.5, when OMP-positive mature OSNs started to appear




Fig. 1. *In Situ* Hybridization of Olfactory Marker Protein and TRPV2 mRNA in Olfactory Mucosa of Adult Mice

Digoxigenin-labeled RNA probes specific to olfactory marker protein (a) and TRPV2 (b) were hybridized with cryosections of olfactory mucosa and visualized with the colorant developed to brown precipitates. OSN, olfactory sensory neurons. The expression pattern shown here was confirmed in two independent experiments. Scale bar, 100  $\mu$ m.




Fig. 2. Accumulations of TRPV2 Protein at Olfactory Axon Bundles

(a–c). Olfactory mucosa was double-labeled with anti-TRPV2 antibody (a) and olfactory marker protein antibody for staining of the mature olfactory neurons (b). TRPV2-immunopositive bundles are indicated by arrows in (a). (c). Overlaid images of TRPV2 (green) and olfactory marker protein (red). (d–f). Double immunolabeling with TRPV2 and growth-associated protein-43, the latter of which is a typical marker of immature olfactory sensory neurons. TRPV2-labeled axons (d, arrows) are co-localized with growth-associated protein-43-immunoreactive axons (e). A merged image of TRPV2 (green) and growth-associated protein-43 (red) is shown in (f). The expression pattern shown here was confirmed in three independent experiments. Scale bar, 100  $\mu$ m.




Fig. 3. Localization of Mouse TRPV2 in Olfactory Bulb

Olfactory bulb sections were stained by double immunofluorescence with olfactory marker protein (a) and TRPV2 antibody (b); a merged image of olfactory marker protein (red) and TRPV2 (green) is shown (c). The expression pattern shown here was confirmed in three independent experiments. Scale bar, 100  $\mu$ m.




Fig. 4. *In Situ* Hybridization of TRPV2 and Insulin-Like Growth Factor-I Receptor mRNA in Olfactory Mucosa of Adult Mice

Digoxigenin-labeled RNA probes specific to TRPV2 (a) and insulin-like growth factor-I receptor (b) were hybridized with cryosections of olfactory mucosa. The expression pattern shown here was confirmed in two independent experiments. Scale bar, 100  $\mu$ m. As shown in Fig. 1, TRPV2 mRNA was expressed in the subpopulation of olfactory neurons (a, green). Insulin-like growth factor-I receptor mRNA was also expressed in the subpopulation of olfactory neurons (b, red). Most neurons expressing TRPV2 mRNA co-expressed insulin-like growth factor-I receptor mRNA. Note that some neurons expressing insulin-like growth factor-I receptor mRNA did not express TRPV2 mRNA.




Fig. 5. Analysis of TRPV2 Expression in Nasal Regions during Fetal Development

(a), Architecture of the nasal region in E12.5 embryonic stages is shown by staining with nuclei. In (a)–(c), dorsal is to the top, frontal is to the bottom, medial is to the right, and lateral is to the left. A rectangle indicates the area shown in panels (b) and (c). FB, forebrain; OE, olfactory epithelium. Double immunofluorescence for  $\beta$ III-tubulin (b, cytoskeletal protein localized in cell bodies, axons, and dendrites of neurons) and TRPV2 (c) at E12.5. Arrows indicate neuroepithelium co-stained with TRPV2 and  $\beta$ III-tubulin. Sagittal section of an E14.5 mouse nose simultaneously stained for  $\beta$ III-tubulin (d), TRPV2 (e). Sagittal section of an E14.5 mouse nose simultaneously stained for olfactory marker protein (f) and TRPV2 (g). (h), The merged images of TRPV2 pseudocolored with green and olfactory marker protein with red are shown. The expression pattern shown here was confirmed in three independent experiments. Scale bar, 100  $\mu$ m. Pictures in b–h were photographed in the same magnification.

(Fig. 5f), axons derived from the OSNs appeared as nerve bundles stained with  $\beta$ III-tubulin between developing OEs, as shown in Fig. 5d. TRPV2 immunoreactivity co-localized at neurons labeled with  $\beta$ III-tubulin, clearly demonstrating that TRPV2 protein localizes on axon bundles of the OSNs during fetal development (Figs. 5d, e). At this stage, staining with TRPV2 antibody is visible on cell bodies and dendrites of olfactory neurons lying in developing OE. Double immunofluorescence using OMP and TRPV2 demonstrated staining of OMP-positive olfactory neurons with TRPV2 antibody (Fig. 5f–h), clearly showing that OMP-positive mature OSNs express TRPV2. Staining for TRPV2 using sagittal sections

from E17.5 embryo was similar to that from E14.5 embryos (not shown).

## DISCUSSION

In this study, we established that TRPV2 mRNA is expressed in a subset of OSNs and that protein is localized on olfactory axons. TRPV2-positive axons innervate to the olfactory nerve layer and glomerular layer in the OB. Localization of TRPV2 was also observed in olfactory axon bundles in embryonic stages, suggesting that TRPV2 has an essential role in processes related to the growth of olfactory axons during

fetal development.

**Restriction of TRPV2 Localization to Olfactory Axon Bundles *in Vivo*** In the adult olfactory system, TRPV2 distribution is restricted to the axons and is hardly observed in the soma and dendrites. Neurite outgrowth is regulated by cytoplasmic calcium-level elevation controlled by calcium-permeable channel activity on plasma membranes.<sup>14)</sup> The members of the TRP family are good candidates for the  $\text{Ca}^{2+}$  influx pathway. For example, TRPC1 is compartmentalized at the growth cone of *Xenopus* spinal neurons, and is required for the turning of growth cones induced by netrin-1 and brain-derived neurotrophic factor.<sup>25-27)</sup> The TRPC5 channel, which is localized to growth cones and gradually disappears as neurons mature and form stable connections with neighboring neurons, is a component of neurite extension by changing the growth cone morphology in rat hippocampal neurons.<sup>28)</sup> These results suggest that the TRP family plays a significant role in regulating the cytoplasmic calcium levels in axon guidance. The immunolocalization of TRPV2 on axon bundles in adult olfactory mucosa and in the GAP43-positive immature axons in the OB suggests that TRPV2 preferentially accumulates in the elongating axons of OSNs in the OE and OB. It is therefore possible that TRPV2 accumulated in elongating OSN axons is a driver of localized increases in intracellular calcium level as a means of extending the axons.

**Putative Functional Involvement of TRPV2 in the Event Controlled by IGF-I in Olfactory Sensory Neurons** IGF-I promotes myelination of neuronal axons, formation of synaptic networks, synaptic plasticity, cognitive function, and the survival and longevity of cells in developing and adult neurons.<sup>29)</sup> Evidence implicates IGF-I in the development and regeneration of neurons in the olfactory system. In the present study, we showed that IGF-IR mRNA exists in a subset of the mouse OSNs as well as in the rat OSNs.<sup>30,31)</sup> In the rat, mRNA encoding IGF-I and IGF-IR exist in the OB.<sup>10,11)</sup> Mucus covering the human OE has also been reported to contain IGF-I.<sup>32)</sup> In cultured olfactory neurons, IGF is a chemoattractant for axon growth cones.<sup>33)</sup> Therefore, it is possible that IGF-I generated in the OB and/or OE activates OSNs *via* the IGF-IR.

Mitral cell layer formation is disrupted in the olfactory bulb of *Igf-1<sup>-/-</sup>* mice.<sup>34)</sup> In chick olfactory mucosa, IGF-I infusion suppresses the apoptotic cell death of OSNs, as well as transient effects on the maintenance of differentiation and mitosis stimulation of basal cells after olfactory axotomy.<sup>35)</sup> In the adult rat, only a few glomeruli were IGF-IR-positive, some of which were unusually small and strongly labeled. Some IGF-IR-positive fibers penetrated deeper into the external plexiform layer, even in adults. In developing tissues, IGF-IR staining co-localized with that for OMP and GAP43.<sup>12)</sup> TRPV2 has been identified as a channel regulated by IGF-I through PI3K and the Akt signaling axis in ectopically expressed mammalian cell lines.<sup>18)</sup> We have demonstrated that the distribution of the IGF-IR mRNA is consistent with the localization of TRPV2 mRNA-expressing neurons in the OE.

IGF-I induced increases in  $[\text{Ca}^{2+}]_i$ , *via* the translocation of TRPV2 from intracellular pools to the plasma membrane in Chinese hamster ovary (CHO) cells expressing TRPV2.<sup>18)</sup> It is possible that IGF-I regulates both the rapid traveling of TRPV2 into axons of olfactory neurons and the local elevation in cytoplasmic calcium levels of olfactory axons and that TRPV2 mediates a mitotic, differentiation, and survival cues

from IGF-I in the OSNs. It is particularly interesting that IGF-I has been shown to promote peripheral axonal regeneration in sensory neurons *in vitro* and *in vivo*.<sup>36,37)</sup> These findings illustrate that TRPV2 is an IGF-I-operated cytoplasmic calcium ion regulatory channel for axonal growth, and is required for the regeneration of sensory neurons expressing TRPV2, although further investigations of the role of TRPV2 against neurite outgrowth *in vivo* and *in vitro* are needed.

**TRPV2 Is Expressed in Developing Olfactory Neurons** TRPV2-expressing cells have been found in dorsal root ganglia during fetal development.<sup>38)</sup> In this study, we investigated TRPV2 expression in the olfactory system from E10.5 to E17.5. TRPV2-immunopositive axons existing from the presumptive OE were detected as early as E12.5. TRPV2 immunopositive neurons were co-labeled with  $\beta$ III-tubulin starting from E12.5 and OMP starting from E14.5, and this showed TRPV2 protein is expressed in immature and mature olfactory neurons, and is localized in axons of the developing OSNs. The early stages of OE development in the mouse have been described at the morphological level.<sup>39,40)</sup> The placodes invaginate, forming the olfactory pits, and OSNs start to grow axons at E10.5. Proliferating OSN progenitors expressing Mash1 and proliferating OSN precursor expressing NeuroD emerge in developing OE at *ca.* E11.5.<sup>41)</sup> The OE becomes organized in layers around E11.5–12.5, with OSN progenitors becoming progressively localized to the basal side of the epithelium and post-mitotic neurons to an intermediate zone.

Olfactory pioneer axons first reach the OB anlagen at the anterior tip of the telencephalon at E12.5<sup>42)</sup> and as shown in Fig. 5b by staining with  $\beta$ III-tubulin in the anterior tip of the forebrain. From E14.5 to E17.5, axodendritic synapses start to form on the dendritic growth cones of mitral cells.<sup>43)</sup> We observed no significant signals in the anterior tips of the embryonic forebrain, prospective OB, at E12.5 and E14.5. Together, the spatiotemporal TRPV2 expression in the embryonic nasal region correlates with the growth and maintenance of olfactory axons but not with the commitment, differentiation, migration, and propagation of olfactory progenitor (Mash1-positive) and precursor (NeuroD-positive) cells, or with the formation of synaptic connection in the developing OB. IGF-IR immunoreactivities are found in the olfactory nerve layer of the rat OB at E18 and 19.<sup>12)</sup> Therefore, it is possible that TRPV2 coupled with IGF-IR also concerns with growing olfactory axons in the OSNs during development as similar to during regeneration. Obviously, further experimental studies are needed to verify this hypothesis.

**Acknowledgment** We deeply appreciate Dr. F. L. Margolis for his gift of the anti-OMP antiserum.

## REFERENCES

- 1) Graziadei PP, Graziadei GA. Neurogenesis and neuron regeneration in the olfactory system of mammals. I. Morphological aspects of differentiation and structural organization of the olfactory sensory neurons. *J. Neurocytol.*, 8, 1–18 (1979).
- 2) Graziadei GA, Graziadei PP. Neurogenesis and neuron regeneration in the olfactory system of mammals. II. Degeneration and reconstitution of the olfactory sensory neurons after axotomy. *J. Neurocytol.*, 8, 197–213 (1979).
- 3) Holcomb JD, Mumm JS, Calof AL. Apoptosis in the neuronal lin-

- age of the mouse olfactory epithelium: Regulation *in vivo* and *in vitro*. *Dev. Biol.*, **172**, 307–323 (1995).
- 4) Mahanthappa NK, Schwarting GA. Peptide growth factor control of olfactory neurogenesis and neuron survival *in vitro*: Roles of EGF and TGF- $\beta$ s. *Neuron*, **10**, 293–305 (1993).
  - 5) Hansel DE, Eipper BA, Ronnett GV. Regulation of olfactory neurogenesis by amidated neuropeptides. *J. Neurosci. Res.*, **66**, 1–7 (2001).
  - 6) Bauer S, Rasika S, Han J, Mauduit C, Raccurt M, Morel G, Jourdan F, Benahmed M, Moysé E, Patterson PH. Leukemia inhibitory factor is a key signal for injury-induced neurogenesis in the adult mouse olfactory epithelium. *J. Neurosci.*, **23**, 1792–1803 (2003).
  - 7) Kawauchi S, Beites CL, Crocker CE, Wu HH, Bonnin A, Murray R, Calof AL. Molecular signals regulating proliferation of stem and progenitor cells in mouse olfactory epithelium. *Dev. Neurosci.*, **26**, 166–180 (2004).
  - 8) Zhang X, Klueber KM, Guo Z, Lu C, Roisen FJ. Adult human olfactory neural progenitors cultured in defined medium. *Exp. Neurol.*, **186**, 112–123 (2004).
  - 9) Beites CL, Kawauchi S, Crocker CE, Calof AL. Identification and molecular regulation of neural stem cells in the olfactory epithelium. *Exp. Cell Res.*, **306**, 309–316 (2005).
  - 10) Rotwein P, Burgess SK, Milbrandt JD, Krause JE. Differential expression of insulin-like growth factor genes in rat central nervous system. *Proc. Natl. Acad. Sci. U.S.A.*, **85**, 265–269 (1988).
  - 11) Yamaguchi F, Itano T, Mizobuchi M, Miyamoto O, Jangua NA, Matsui H, Tokuda M, Ohmoto T, Hosokawa K, Hatase O. Insulin-like growth factor I (IGF-I) distribution in the tissue and extracellular compartment in different regions of rat brain. *Brain Res.*, **533**, 344–347 (1990).
  - 12) Ferrari CC, Johnson BA, Leon M, Pixley SK. Spatiotemporal distribution of the insulin-like growth factor receptor in the rat olfactory bulb. *Neurochem. Res.*, **28**, 29–43 (2003).
  - 13) Zhang C, Finger TE, Restrepo D. Mature olfactory receptor neurons express connexin 43. *J. Comp. Neurol.*, **426**, 1–12 (2000).
  - 14) Henley J, Poo MM. Guiding neuronal growth cones using Ca<sup>2+</sup> signals. *Trends Cell Biol.*, **14**, 320–330 (2004).
  - 15) Clapham DE. TRP channels as cellular sensors. *Nature*, **426**, 517–524 (2003).
  - 16) Caterina MJ, Rosen TA, Tominaga M, Brake AJ, Julius D. A capsaicin-receptor homologue with a high threshold for noxious heat. *Nature*, **398**, 436–441 (1999).
  - 17) Stein AT, Ufret-Vincenty CA, Hua L, Santana LF, Gordon SE. Phosphoinositide 3-kinase binds to TRPV1 and mediates NGF-stimulated TRPV1 trafficking to the plasma membrane. *J. Gen. Physiol.*, **128**, 509–522 (2006).
  - 18) Ahmed MK, Takumida M, Ishibashi T, Hamamoto T, Hirakawa K. Expression of transient receptor potential vanilloid (TRPV) families 1, 2, 3 and 4 in the mouse olfactory epithelium. *Rhinology*, **47**, 242–247 (2009).
  - 19) Nakashimo Y, Takumida M, Fukui T, Anniko M, Hirakawa K. Expression of transient receptor potential channel vanilloid (TRPV) 1–4, melastin (TRPM) 5 and 8, and ankyrin (TRPA1) in the normal and methimazole-treated mouse olfactory epithelium. *Acta Otolaryngol.*, **130**, 1278–1286 (2010).
  - 20) Kanzaki M, Zhang YQ, Mashima H, Li L, Shibata H, Kojima I. Translocation of a calcium-permeable cation channel induced by insulin-like growth factor-I. *Nat. Cell Biol.*, **1**, 165–170 (1999).
  - 21) Keller A, Margolis FL. Immunological studies of the rat olfactory marker protein. *J. Neurochem.*, **24**, 1101–1106 (1975).
  - 22) Schaeren-Wiemers N, Gerfin-Moser A. A single protocol to detect transcripts of various types and expression levels in neural tissue and cultured cells: *in situ* hybridization using digoxigenin-labelled cRNA probes. *Histochemistry*, **100**, 431–440 (1993).
  - 23) Verhaagen J, Oestreicher AB, Gispén WH, Margolis FL. The expression of the growth associated protein B50/GAP43 in the olfactory system of neonatal and adult rats. *J. Neurosci.*, **9**, 683–691 (1989).
  - 24) Roskams AJ, Bethel MA, Hurt KJ, Ronnett GV. Sequential expression of Trks A, B, and C in the regenerating olfactory neuroepithelium. *J. Neurosci.*, **16**, 1294–1307 (1996).
  - 25) Wang GX, Poo MM. Requirement of TRPC channels in netrin-1-induced chemotropic turning of nerve growth cones. *Nature*, **434**, 898–904 (2005).
  - 26) Li Y, Jia YC, Cui K, Li N, Zheng ZY, Wang YZ, Yuan XB. Essential role of TRPC channels in the guidance of nerve growth cones by brain-derived neurotrophic factor. *Nature*, **434**, 894–898 (2005).
  - 27) Shim S, Goh EL, Ge S, Sailor K, Yuan JP, Roderick HL, Bootman MD, Worley PF, Song H, Ming GL. XTRPC1-dependent chemotropic guidance of neuronal growth cones. *Nat. Neurosci.*, **8**, 730–735 (2005).
  - 28) Greka A, Navarro B, Oancea E, Duggan A, Clapham DE. TRPC5 is a regulator of hippocampal neurite length and growth cone morphology. *Nat. Neurosci.*, **6**, 837–845 (2003).
  - 29) Wada A, Yokoo H, Yanagita T, Kobayashi H. New twist on neuronal insulin receptor signaling in health, disease, and therapeutics. *J. Pharmacol. Sci.*, **99**, 128–143 (2005).
  - 30) Pixley SK, Dangoria NS, Odoms KK, Hastings L. Effects of insulin-like growth factor I on olfactory neurogenesis *in vivo* and *in vitro*. *Ann. N. Y. Acad. Sci.*, **855**, 244–247 (1998).
  - 31) Suzuki Y, Takeda M. Expression of insulin-like growth factor family in the rat olfactory epithelium. *Anat. Embryol. (Berl.)*, **205**, 401–405 (2002).
  - 32) Federico G, Maremmani C, Cinquanta L, Baroncelli GI, Fattori B, Saggese G. Mucus of the human olfactory epithelium contains the insulin-like growth factor-I system which is altered in some neurodegenerative diseases. *Brain Res.*, **835**, 306–314 (1999).
  - 33) Scolnick JA, Cui K, Duggan CD, Xuan S, Yuan XB, Efstratiadis A, Ngai J. Role of IGF signaling in olfactory sensory map formation and axon guidance. *Neuron*, **57**, 847–857 (2008).
  - 34) Vicario-Abejón C, Yusta-Boyo MJ, Fernandez-Moreno C, de Pablo F. Locally born olfactory bulb stem cells proliferate in response to insulin-related factors and require endogenous insulin-like growth factor-I for differentiation into neurons and glia. *J. Neurosci.*, **23**, 895–906 (2003).
  - 35) Mathonnet M, Comte I, Lalloue F, Ayer-Le Lievre C. Insulin-like growth factor I induced survival of axotomized olfactory neurons in the chick. *Neurosci. Lett.*, **308**, 67–70 (2001).
  - 36) Fernyhough P, Willars GB, Lindsay RM, Tomlinson DR. Insulin and insulin-like growth factor I enhance regeneration in cultured adult rat sensory neurons. *Brain Res.*, **607**, 117–124 (1993).
  - 37) Gutiérrez-Ospina G, Calikoglu AS, Ye P, D'Ercole AJ. *In vivo* effects of insulin-like growth factor-I on the development of sensory pathways: analysis of the primary somatic sensory cortex (S1) of transgenic mice. *Endocrinology*, **137**, 5484–5492 (1996).
  - 38) Tamura S, Morikawa Y, Senba E. TRPV2, a capsaicin receptor homologue, is expressed predominantly in the neurotrophin-3-dependent subpopulation of primary sensory neurons. *Neuroscience*, **130**, 223–228 (2005).
  - 39) Cuschieri A, Bannister LH. The development of the olfactory mucosa in the mouse: light microscopy. *J. Anat.*, **119**, 277–286 (1975).
  - 40) Cuschieri A, Bannister LH. The development of the olfactory mucosa in the mouse: electron microscopy. *J. Anat.*, **119**, 471–498 (1975).
  - 41) Cau E, Casarosa S, Guillemot F, Mashl and Ngn1 control distinct steps of determination and differentiation in the olfactory sensory neuron lineage. *Development*, **129**, 1871–1880 (2002).
  - 42) Yoshihara S, Omichi K, Yanazawa M, Kitamura K, Yoshihara Y. Arx homeobox gene is essential for development of mouse olfactory system. *Development*, **132**, 751–762 (2005).
  - 43) Brunjes PC, Frazier LL. Maturation and plasticity in the olfactory system of vertebrates. *Brain Res.*, **396**, 1–45 (1986).