

AMCoR

Asahikawa Medical University Repository <http://amcor.asahikawa-med.ac.jp/>

Investigative Ophthalmology and Visual Science (2016.9) 57(11):4962–4969.

Glial Endothelin-1 Regulates Retinal Blood Flow During Hyperoxia in Cats

Youngseok Song, Taiji Nagaoka, Takafumi Yoshioka, Shinji Ono, Takanari Wada, Seigo Nakabayashi, Tomofumi Tani, Akitoshi Yoshida

Glial Endothelin-1 Regulates Retinal Blood Flow During Hyperoxia in Cats

Youngseok Song, Taiji Nagaoka, Takafumi Yoshioka, Shinji Ono, Takanari Wada, Seigo Nakabayashi, Tomofumi Tani, and Akitoshi Yoshida

Department of Ophthalmology, Asahikawa Medical University, Asahikawa, Japan

Correspondence: Taiji Nagaoka, Department of Ophthalmology, Asahikawa Medical University, Midorigaoka Higashi 2-1-1, Asahikawa, 078-8510, Japan; nagaoka@asahikawa-med.ac.jp.

Submitted: March 20, 2016

Accepted: July 17, 2016

Citation: Song Y, Nagaoka T, Yoshioka T, et al. Glial endothelin-1 regulates retinal blood flow during hyperoxia in cats. *Invest Ophthalmol Vis Sci*. 2016;57:4962-4969. DOI:10.1167/iov.16-19599

PURPOSE. To investigate the role of endothelin-1 (ET-1) in retinal glial cells in regulating retinal blood flow (RBF) during hyperoxia in cats.

METHODS. We measured the vessel diameter (D), blood velocity (V), and blood flow (F) simultaneously in first-order retinal arterioles using a laser Doppler velocimetry system. The animals were under general anesthesia during hyperoxia (100% oxygen) for 10 minutes 24 hours after intravitreal injection of L-2-aminoadipic acid (LAA), a gliotoxic compound, or diluted hydrochloric acid (0.01 N) used as the vehicle control. We also measured the changes in the RBF after intravitreal injection of BQ-123, a specific ET type A receptor antagonist, in LAA-treated eyes. To examine if endothelin-converting enzyme-1 (ECE-1), as an ET-1-generating enzyme located in retinal glial cells, immunohistochemical examinations with costaining of antiglial fibrillary acidic protein (GFAP) antibody and anti-ECE-1 antibody were performed in whole-mount retinas.

RESULTS. During hyperoxia, the decreases in D, V, and F in response to hyperoxia were attenuated significantly ($P < 0.01$ for all comparisons) in the LAA-treated eyes compared with the vehicle control (LAA, D, $-8.5 \pm 1.5\%$; V, $-13.8 \pm 1.5\%$; F, $-27.8 \pm 3.0\%$ versus vehicle control, D, $-16.8 \pm 1.3\%$; V, $-26.3 \pm 2.0\%$; F, $-48.9 \pm 2.4\%$). In LAA-treated eyes, intravitreal injections of BQ-123 did not change the rate of hyperoxia-induced RBF compared to LAA-treated eyes. The anti-ECE-1 antibody was costained with anti-GFAP antibody in the whole-mount retinas.

CONCLUSIONS. The current findings suggest that retinal glial ET-1 may play an important role in regulating RBF during hyperoxia in cats.

Keywords: retinal blood flow, retinal glial cells, hyperoxia, endothelin, endothelin converting enzyme

Glial cells, including Müller cells and astrocytes as the main glial cells in the retina,¹ are vital for maintaining normal retinal function.² Recent animal experiments have shown that glial cells play a principal role in regulating retinal blood flow (RBF) in various conditions, such as increased intraocular pressure (IOP) and photic stimulation.^{3,4} We also showed previously that glial cells play a crucial role in regulating RBF in flicker-induced hyperemia in cats.⁵ Indeed, impaired glial cellular activity might be related to the pathologic mechanisms of ocular disorders such as diabetic retinopathy (DR) and glaucoma.^{6,7} Moreover, some clinical studies have reported that the hyperoxia-induced vasoconstriction in the retina deteriorates in patients with these diseases.^{8,9} However, it remains unclear how glial cells regulate RBF during hyperoxia.

Systemic hyperoxia constricts the retinal arterioles and decreases the RBF in humans and animals through release of endothelin (ET)-1.¹⁰⁻¹⁴ Previously, we also reported that ET-1 plays a major role in hyperoxia-induced vasoconstriction in cats.¹⁵ Endothelin-1 is a 21 amino acid polypeptide, expressed in both endothelial and glial components in the retina,¹⁶ that has a potent constrictor action by binding to the high-affinity ET type A (ETA) receptor in retinal vascular smooth muscle cells and pericytes.¹⁷⁻²¹ Endothelin-1 is generated from proendothelin-1, the precursor of ET-1, via the action of endothelin-converting enzyme-1 (ECE-1).^{22,23} Takagi et al.¹²

reported that administration of exogenous ET-1 was associated with decreased RBF in rats. Surprisingly, those investigators also reported that ET-1 did not increase significantly in cultured bovine retinal endothelial cells exposed to hyperoxia in Northern blot analysis,¹² suggesting that vascular endothelial cells might not be involved with production of ET-1 in response to systemic hyperoxia. These findings taken together, we speculated that glial cells might have a role in regulating RBF during hyperoxia via production of ET-1.

We showed previously that intravitreal injection of LAA, a gliotoxic compound, had a specific role in glial cells,⁵ and BQ-123, a specific ETA antagonist, had a specific role in the ETA receptor in the feline retina.¹⁵ In the current study, we hypothesized that ET-1 in glial cells might have a role in regulating RBF during hyperoxia after suppressing the retinal glial cellular function and antagonizing the ETA receptor with intravitreal injection of LAA and BQ-123 in cats.

MATERIALS AND METHODS

Animal Preparation

The Animal Care Committee of Asahikawa Medical University approved the protocols for the use of animals, which adhered to the ARVO Statement for the Use of Animals in Ophthalmic and



Vision Research. Fifteen adult European shorthair cats provided by Shiraishi Laboratory Animals Co., Ltd. (3.4–4.5 kg; Koshigaya, Saitama, Japan) of either sex were tracheostomized and ventilated mechanically with room air containing 2.0% sevoflurane. The flow rate of sevoflurane was maintained at 1.5 L/min during the experiment. Catheters were placed in the femoral arteries and vein. The mean arterial blood pressure (MABP) and heart rate (HR) were monitored continuously with a transducer (PowerLab; ADInstruments, Inc., Colorado Springs, CO, USA) and recorder (LabChart; ADInstruments, Inc.) in the proximal thoracic descending aorta. Pancuronium bromide (0.1 mg/kg/h) (Daichi Sankyo Co., Tokyo, Japan) was infused continuously via the femoral vein to maintain skeletal muscle relaxation during the experiment. With the animal prone, the head was fixed in a stereotaxic instrument. Arterial pH, arterial partial carbon dioxide tension (PaCO_2), arterial partial oxygen tension (PaO_2), and bicarbonate ion (HCO_3^-) were measured intermittently with a blood gas analyzer (model ABL5; Radiometer, Copenhagen, Denmark). The rectal temperature was measured and maintained between 37°C and 38°C with a heated blanket. The pupils were dilated with 0.5% tropicamide (Santen Pharmaceutical Co., Osaka, Japan). A 0-diopter contact lens (Seed Co. Ltd., Tokyo, Japan) was placed on the cornea, which was protected by instillation of a drop of sodium hyaluronate (Healon; Abbott Medical Optics, Inc., Abbott Park, IL, USA). A 26-gauge butterfly needle was inserted into the anterior chamber and connected to a pressure transducer and a balanced salt solution (Alcon, Fort Worth, TX, USA) reservoir for monitoring and maintaining the IOP at 10 mm Hg, respectively.

RBF Measurements

A laser Doppler velocimetry system (Laser Blood Flowmeter, model 100; Canon, Inc., Tokyo, Japan) customized for felines was used to measure the retinal arteriolar diameter (D) (in micrometers) and velocity (V) (mm/s) as described previously.^{24,25} The RBF in the arterioles ($\mu\text{L}/\text{min}$) was calculated based on the acquired V and D. Laser Doppler measurements of the temporal retinal arterioles were performed in one eye of each animal. We studied the first-order arterioles because they have relatively straight segments and were sufficiently distant from the adjacent vessels for consistent measurements.

The RBF was calculated using the formula $RBF = S \times V_{mean}$, where S is the cross-sectional area of the retinal arteriole at the laser Doppler measurement site, assuming a circular cross section, and V_{mean} is the mean blood V calculated as $V_{mean} = V_{max} / 2$.²⁶ The MABP was determined using the formula $MABP = \text{diastolic BP} + (\text{systolic BP} - \text{diastolic BP})/3$, which is the index of the systemic BP. Because the cats were prone during the experiments, the ocular perfusion pressure (OPP) was calculated as $OPP = MABP - IOP$.^{27,28}

Intravitreal Injections and Chemicals

A 30-gauge needle (100- μL syringe; Hamilton, Reno, NV, USA) was used to perform the intravitreal injections 3 mm posterior to the limbus with care taken to not injure the lens and retina.²⁴ The head of the needle was positioned over the optic disc region. All drugs were purchased from Sigma-Aldrich Corp. (St. Louis, MO, USA). BQ-123 was dissolved in phosphate-buffered saline (PBS). L-2-aminoadipic acid was dissolved in 0.01 N hydrochloric acid (HCl) because LAA does not dissolve in PBS.⁵ The volume of the intravitreal injections was 50 μL , which does not alter any retinal circulatory parameters and minimizes the systemic effects of the inhibitors.²⁴ Because the volume of the cat vitreous is approximately 2.5 mL, the solution (50 μL) injected into the vitreous cavity is diluted by a factor of 50 near the retinal

vessels. Hereafter, we refer to drug concentrations as injected concentrations.

The 60 mM concentration of LAA (estimated final concentrations in the vitreous cavity were 1.2 mM) was chosen because 1.25 mM DL- α -aminoadipic acid causes swelling of the Müller cells and astrocytes and the remaining neural cells are intact.^{5,29} As we reported previously,⁵ an observation period of 24 hours after intravitreal injection of LAA was used because no pathologic changes or dysfunction on electroretinography (ERG) was observed in the neural retina and the maximal responses in RBF were obtained in our previous experiment.³

Induction of Hyperoxia

Hyperoxia was induced by inhalation of 100% oxygen for 10 minutes 24 hours after injection of 0.01 N HCl or LAA into each cat. The RBF was measured every minute during and after hyperoxia. An average of five measurements obtained at 2-minute intervals was defined as the baseline value before initiation of hyperoxia (Fig. 1). Blood gas analysis was performed before, at the end, and 10 minutes after the end of hyperoxia.

Effect of Intravitreal Injection of BQ-123 After Initiation of Hyperoxia in LAA-Treated Eyes

We used BQ-123 to study if the suppressed responses in RBF during hyperoxia in LAA-treated eyes resulted from the action of ET. BQ-123, an ETA receptor antagonist, was injected into the vitreous for an extracellular concentration of 2.0×10^{-5} M near the retinal vessels. This concentration ensures a maximal retinal hemodynamic response.¹³ Hyperoxia was induced 60 minutes after injection of BQ-123 because we confirmed previously that the effects of BQ-123 (1 mM) reached the maximal level 60 minutes after the injection.¹⁵ BQ-123 was injected 23 hours after LAA to confirm the maximal responses of LAA and BQ-123 (Fig. 1). The RBF was measured every minute during and after hyperoxia (Fig. 1).

Immunohistochemistry

For whole-mount assessment, the eyes were enucleated and fixed in 1% paraformaldehyde for 1 hour. The retina was blocked and permeabilized in 5% goat serum with 0.3% Triton (Wako Pure Chemical Industries, Ltd., Osaka, Japan) in PBS for 1 to 2 hours. The retinas were transferred to primary antibodies diluted in block solution and incubated for 1 hour at room temperature. We used the following specific primary antibodies: a mouse anti-glial fibrillary acidic protein (GFAP)-cy3 antibody (1:400; Sigma-Aldrich Corp.) and a goat anti-ECE-1 (1:200; R & D Systems, Minneapolis, MN, USA). The secondary antibody was anti-goat IgG Alexa Fluor 647-conjugated antibody (Abcam, Inc., Cambridge, MA, USA). The slides were mounted (Dako, Tokyo, Japan), observed for green (cy3) and red (Alexa Fluor 647) staining, and analyzed with a fluorescence microscope (FluoView FV1000; Olympus, Tokyo, Japan). The emission wavelengths were 567 nm (575- to 620-nm band-pass filter, cy3) and 647 nm (655- to 755-nm band-pass filter, Alexa Fluor 647).

Ex Vivo Experiment Using Isolated Retinal Arterioles

We used the porcine retinal arterioles to confirm the effects of LAA on the contractility of the retinal arterioles in cats. The eyes were enucleated immediately from pigs of either sex (age, 16–24 weeks; weight, 25–35 kg) after the animals were killed




FIGURE 1. The scheme of the time interval and procedures in the RBF. Hyperoxia was induced by inhalation of 100% oxygen for 10 minutes 24 hours after injection of 0.01 N HCl (vehicle control) or LAA at 60 mM into the vitreous cavity because the effects of LAA (60 mM) reached the maximal level 24 hours after the injection. The RBF was measured every minute during and after hyperoxia (10 minutes). An average of five measurements obtained at 2-minute intervals was defined as the baseline value before initiation of hyperoxia. In addition, BQ-123 was injected 23 hours after injection of 0.01 N HCl (vehicle control) or LAA at 60 mM since the effects of BQ-123 (1 mM) reached the maximal level 60 minutes after the injection. The eyes were enucleated for whole-mount assessment after RBF measurements. 23 hours, 23 hours after injection of LAA (60 mM) or 0.01 N HCl; 24 hours, 24 hours after injection of LAA (60 mM) or 0.01 N HCl.

in a local abattoir and transported to the laboratory in a moist chamber on ice.

Isolation and Cannulation of Microvessels

The techniques used to identify, isolate, cannulate, pressurize, and visualize the retinal microvessels have been described previously.^{30–33} Briefly, single second-order retinal arterioles (90–110 μm in situ) were dissected with microdissection forceps and the isolated retinal arterioles were cannulated with a pair of glass micropipettes and pressurized to 55-cm H₂O intraluminal pressure without flow using two independent pressure reservoir systems.³⁴ The internal diameter of the isolated vessels was recorded continuously using video microscopic techniques throughout the experiments.³⁵

Control Experiment

Cannulated and pressurized arterioles were bathed in physiological saline solution with albumin (0.1%) at 36°C to 37°C to allow development of basal tone. After the vessels developed a stable basal tone (~30–40 minutes), they were incubated with LAA 60 mM or 0.01 N HCl (vehicle control) for an hour. Thereafter, the vessels were exposed to ET-1 at 1 nM (Sigma-Aldrich Corp.), and the ET-1-induced vasoconstriction was evaluated every 5 minutes for 20 minutes. After 20 minutes, we regarded the vessel caliber as the maximal response.

Statistical Analysis

All data are expressed as the mean percentage ± standard error of the mean. The changes in RBF were calculated as percentage decreases from the baseline measurements. For statistical analysis, we used analysis of variance (ANOVA) for repeated measurements, followed by post hoc comparisons with the Dunnett procedure. Group comparisons of the RBF, systemic parameters, and the ET-1-induced vasoconstrictions in isolated retinal arterioles were performed using the Mann-Whitney *U* test or Wilcoxon signed rank test. *P* < 0.05 was considered significant.

RESULTS

Systemic Changes in Response to Hyperoxia

During hyperoxia, there were no significant differences in the pH, PaCO₂, HCO₃⁻, MABP, HR, or OPP, whereas the PaO₂ increased significantly (*P* < 0.01) in all groups (Table 1). Intravitreal injection of BQ-123 did not alter any systemic parameters (Supplementary Table S1).

Effects of LAA and BQ-123 on RBF at Rest

Twenty-four hours after injection of LAA (60 mM) alone or LAA (60 mM) + BQ-123 (1 mM), there were no significant changes in the retinal circulatory parameters (*n* = 5) (Table 2). We preliminarily confirmed the absence of significant differences in the ocular and systemic circulatory parameters at rest between PBS- and the 0.01 N HCl-treated eyes (Supplementary Tables S2, S3).

Effects of LAA on Hyperoxia-Induced Changes in Retinal Circulation

The D, V, and F decreased gradually after induction of hyperoxia compared with baseline in the vehicle control group (Fig. 2A). Ten minutes after the onset of hyperoxia in the vehicle control group, the percentage decreases from baseline were -16.8 ± 1.3% in D, -26.3 ± 2.0% in V, and -48.9 ± 2.4% in F (Fig. 2B). However, 10 minutes after the onset of hyperoxia in LAA-treated eyes, the decreases in the RBF during hyperoxia were attenuated significantly compared with the vehicle controls (LAA 60 mM, D, -8.5 ± 1.5%; V, -13.8 ± 1.5%; F, -27.8 ± 3.0%) (Fig. 2B). After hyperoxia, the retinal circulatory parameters began to return to the baseline values. Ten minutes after the end of hyperoxia, the retinal circulatory parameters recovered to the baseline value in the vehicle controls and LAA-treated eyes. In a preliminary study, we confirmed that there were no significant differences in the ocular and systemic circulatory parameters during hyperoxia

TABLE 1. Changes in Systemic Parameters During Hyperoxia

	0.01 N HCL, n = 5		LAA 60 mM, n = 5		LAA 60 mM + BQ-123 1 mM, n = 5	
	Before	Hyperoxia	After	Before	Hyperoxia	After
pH	7.41 ± 0.03	7.42 ± 0.02	7.42 ± 0.01	7.41 ± 0.02	7.41 ± 0.01	7.41 ± 0.03
PaCO ₂ , mm Hg	29.8 ± 0.7	29.2 ± 0.7	30.4 ± 0.7	30.8 ± 0.4	29.8 ± 0.4	30.1 ± 0.7
PaO ₂ , mm Hg	100.4 ± 3.6	607.6 ± 7.8*	109.0 ± 2.9	109.0 ± 2.5	618.4 ± 10.0*	110.4 ± 5.5
HCO ₃ ⁻ , M	20.5 ± 0.5	20.6 ± 0.4	20.9 ± 0.5	19.8 ± 0.4	19.6 ± 0.3	20.2 ± 0.4
MABP, mm Hg	101.8 ± 2.0	101.3 ± 2.9	100.9 ± 2.4	102.5 ± 0.8	102.5 ± 0.8	103.5 ± 1.9
HR, beats/min	123.8 ± 4.6	124.6 ± 4.8	125.4 ± 5.5	130.2 ± 4.2	129.8 ± 4.2	131.0 ± 4.8
OPP, mm Hg	90.2 ± 2.1	89.7 ± 2.0	89.3 ± 2.5	89.8 ± 1.0	89.8 ± 0.8	91.7 ± 0.2

The data are expressed as the means ± standard errors of the mean. For the comparison before with hyperoxia, we used the Wilcoxon signed rank test. * $P < 0.05$ is considered significant. Before, before induction of hyperoxia; hyperoxia, at the end of hyperoxia; after, 10 minutes after the end of hyperoxia.

TABLE 2. Effect of BQ-123 on Ocular Parameters (n = 5)

	0.01 N HCL	LAA 60 mM	
		LAA 60 mM	+ BQ-123 1 mM
Diameter, μ m	118.1 ± 2.6	115.2 ± 3.1	113.6 ± 4.5
Velocity, mm/s	35.6 ± 2.3	35.0 ± 2.1	33.1 ± 3.1
Flow, μ l/min	11.7 ± 0.8	10.9 ± 0.6	10.5 ± 1.6

The data are expressed as the means ± standard errors of the mean. The Mann-Whitney U test was used to compare the vehicle controls (0.01 N HCL) with LAA or LAA + BQ-123. There are no significant differences in D, V, or F between the groups.

between PBS- and the 0.01 N HCL-treated eyes (Supplementary Fig. S1; Supplementary Table S2).

Effects of BQ-123 in LAA-Treated Eyes During Hyperoxia

As reported previously, 60 minutes after injection of BQ-123 in LAA-treated eyes (before induction of hyperoxia), there were no significant differences in D, V, and F compared with the preinjection levels.

In the LAA (60 mM)-treated eyes with injections of BQ-123 (1 mM), the percentage reductions of all retinal circulatory parameters were comparable to those in the LAA-treated eyes 10 minutes after induction of hyperoxia (Fig. 3). After hyperoxia, the D, V, and F immediately returned to the baseline level (data not shown).

Immunohistochemistry

To examine if the retinal glial cells have endothelin converting enzyme-1 (ECE-1), the costaining of GFAP and ECE-1 was assessed in the whole retina (Fig. 4). Glial fibrillary acidic protein immunofluorescence histochemistry (Fig. 4A), binding of ECE-1 (Fig. 4B), and merged (Fig. 4C) were performed on flat-mounted feline retinal preparations. In the feline retina, ECE was expressed in the retinal glial cells and vessels, although the neurons were not stained (Fig. 4C). In a preliminary study, the control sections were incubated with no primary antibody, with only secondary antibody conjugates, and these sections did not stain (data not shown).

Constriction of LAA-Incubated Isolated Retinal Arterioles Induced by ET-1

We examined the effects of LAA on the contractility resulting from ET-1 (1 nM) in the ex vivo porcine retinal isolated vessels using isolated porcine retinal arterioles. There was no significant difference ($-63.6 \pm 4.7\%$ vs. $-71.9 \pm 2.5\%$; $n = 4$, $P = 0.34$) in the contractility of the retinal arterioles between the LAA 60 mM and 0.01 N HCL groups (Fig. 5).

DISCUSSION

In the current study, we found that decreases in RBF during hyperoxia were suppressed in LAA-treated eyes compared with vehicle control eyes (Fig. 2), suggesting that retinal glial cells might participate in the regulation in RBF during hyperoxia. Previously, Takagi et al.¹² reported that ET-1 did not increase significantly in cultured bovine retinal endothelial cells exposed to hyperoxia in their Northern blot analysis, suggesting that it is reasonable to consider that vascular endothelial cells might not be involved with production of ET-1 in response to systemic hyperoxia. In addition, the




FIGURE 2. Time course of the changes from baseline in the retinal circulation in response to hyperoxia induced by 0.01 N HCl ($n = 5$) as a vehicle control and the LAA 60 mM ($n = 5$) groups. (A) The black bar represents the period of hyperoxia (10 minutes). The data are expressed as mean percentages \pm standard errors (SE) of the means of the baseline values. * $P < 0.05$ and † $P < 0.05$ compared with baseline by 2-way repeated-measures ANOVA followed by the Dunnett procedure. (B) The maximal changes from baseline (10 minutes after




FIGURE 3. The maximal changes from baseline (10 minutes after induction of hyperoxia) comparing LAA-treated eyes with double-blocked eyes (LAA 60 mM + BQ-123 1 mM) in the retinal circulation in response to hyperoxia. The data are expressed as the mean percentages \pm standard errors of the means of the baseline values. * $P < 0.05$ compared with a vehicle control group by 1-way factorial ANOVA followed by the Dunnett procedure. There are no significant differences between the groups.

inhibition ratio of the hyperoxia-induced decrease in RBF in eyes injected with LAA and BQ-123 was comparable to that in eyes treated only with LAA (Fig. 4), suggesting that ET-1 might originate from the retinal glial cells during hyperoxia. This possibility also was supported by immunohistochemistry findings, in which the glial cells and ECE-1 were costained (Fig. 4).

Another possibility is that hyperoxia might increase ET-1 production not only in the glial cells but also in the retinal neurons and trigger an alternative pathway for retinal vessel constriction. Although we could not exclude this possibility, our immunohistochemistry findings also showed that the neurons were not stained by anti-ECE-1 in the feline retina (Fig. 4), suggesting that retinal neurons might not be involved in regulating RBF during hyperoxia via production of ET-1.

A hyperoxia-induced decrease in RBF was not suppressed completely by LAA and/or BQ-123 in the current study (Figs. 2, 3), which suggested that there are other mechanisms besides the glial cells or ET-1 in the reduction of RBF during hyperoxia. Rubanyi and Vanhoutte³⁵ suggested the possible explanation for the residual response in the hyperoxia-induced decrease in RBF that superoxide anions, which were generated from oxygen in hyperoxia, transiently inactivated nitric oxide (NO) as an endothelium-derived relaxing factor and the inactivated NO causes the vasoconstriction in the canine coronary artery.³⁵

We reported previously that L-NG-nitroarginine methyl ester, a NO synthase inhibitor, did not increase the rate by which RBF decreased during hyperoxia,¹⁵ which supported the theory of Rubanyi and Vanhoutte.³⁵ Although we did not examine the effects of superoxide anions on the current results, the transient inactivation of NO by superoxide anions might lead to the residual response besides ET-1 in the hyperoxia-induced decrease in RBF.

In the current study, immunohistochemistry findings indicated that anti-ECE-1 and anti-GFAP were costained in the feline retina (Fig. 4), suggesting that ECE-1 was expressed in the retinal glial cells. In contrast, there was no staining of anti-ET-1 antibody in the glial cells (data not shown), possibly because ET-1 is merely released under normoxic conditions.³⁶ Unfortunately, we could not immediately enucleate eyes under

induction of hyperoxia) in the retinal circulation in response to hyperoxia in the groups in (A). The data are expressed as the mean percentages \pm SEs of the baseline values. * $P < 0.05$ compared with a vehicle control group by 1-way factorial ANOVA followed by the Dunnett procedure.




FIGURE 4. Immunohistochemistry of flat-mounted feline retinas. (A) Anti-GFAP immunofluorescence histochemistry, (B) anti-ECE-1, and (C) merged were performed. ECE is expressed in the feline retinas. Scale: 20 μ m.

the hyperoxic condition because of a technical difficulty in the current study. Alternatively, further experiments using cultured Müller glial cells or retinal astrocytes exposed to hyperoxic conditions are needed to elucidate the mechanisms underlying the hyperoxia-induced increase in ET-1 production in retinal glial cells.

Some ocular diseases, such as DR and glaucoma, impair the regulation of ocular blood flow^{37,38} and reduce the response in hyperoxia-induced vasoconstriction in the retina.^{8,9} In addition, impaired glial cell activity might be related to the pathologic mechanisms of those ocular disorders.³⁹⁻⁴² In the current study, we showed the dysregulation in the RBF during hyperoxia in LAA-treated eyes (Fig. 2). Although we did not use the disease model in this study, the results indicated that glial

dysfunction might be associated with the pathogenesis of DR and/or glaucoma via impaired RBF regulation. Further basic studies using the disease model and more clinical investigations are warranted to examine whether improved glial function is a novel target for treating ocular vascular disorders.

The current study had some limitations. First, the current results did not show which retinal glial cells, the Müller glial cells or astrocytes, play a central role in regulating RBF during hyperoxia, because it is difficult to suppress each cellular function separately pharmacologically using a gliotoxic compound, which affects both of them.^{43,44} Second, although the current data did not provide a definitive explanation for the effect of general anesthesia, we found in a preliminary study that sevoflurane per se did not change the vessel D of isolated




FIGURE 5. Effect of LAA on the contractility of ex vivo porcine retinal isolated arterioles induced by ET-1 (1 nM). The vessels were preincubated with LAA 60 mM ($n = 4$) or 0.01 N HCl (vehicle control, $n = 4$) for an hour. After the preincubation, the vessels were exposed to ET-1 (1 nM), and the maximal responses of the ET-1-induced vasoconstriction were evaluated 20 minutes after exposure to ET-1. There is no significant difference in the contractility of the retinal isolated arterioles between the groups.

porcine retinal arterioles (data not shown), and changes in the concentrations of pancuronium bromide did not alter the RBF in cats anesthetized with sevoflurane (data not shown).⁴⁵ Finally, as we reported previously,⁵ we could not quantify the degree of functional damage in the retinal glial cells after intravitreal injection of LAA at a concentration of 60 mM, although LAA at 60 mM selectively damaged the glial cells without hurting the neurons and retinal vasculature in the feline retinas. More advanced techniques and research are needed to resolve these issues. In the current study, we examined the effects of LAA on the contractility caused by ET-1 (1 nM) in the retinal vessels using isolated porcine retinal arterioles (Fig. 5) and found that there was no significant difference in the contractility of the retinal arterioles between the LAA and the vehicle control groups, suggesting that this compound did not directly affect the contractility of the retinal blood vessels.

In summary, we found that the hyperoxia-induced decrease in RBF was suppressed significantly in LAA-treated eyes and the suppressed response was comparable to that in LAA-treated eyes with intravitreal injection of BQ-123 as the ETA receptor antagonist, suggesting that the glial cells might be involved in regulating RBF during hyperoxia by releasing ET-1 in the glial cells. Because previous reports have indicated the possible involvement of glial cell dysfunction in the pathogenesis of DR and glaucoma,^{6,7} clarifying the detailed mechanisms of the glial cells in regulating the retinal microcirculation might provide further understanding of the pathogenesis in ocular vascular disorders, that is, DR and glaucoma.

Acknowledgments

The authors thank Manami Kuze of Mie University Graduate School of Medicine for her invaluable discussion.

Supported by a Grant-in-Aid for Scientific Research (B) 26861430 from the Ministry of Education, Science, and Culture, Tokyo, Japan (to YS).

Disclosure: **Y. Song**, None; **T. Nagaoka**, None; **T. Yoshioka**, None; **S. Ono**, None; **T. Wada**, None; **S. Nakabayashi**, None; **T. Tani**, None; **A. Yoshida**, None

References

- Hernandez MR. The optic nerve head in glaucoma: role of astrocytes in tissue remodeling. *Prog Retin Eye Res.* 2000;19:297-321.
- Bringmann A, Pannicke T, Grosche J, et al. Müller cells in the healthy and diseased retina. *Prog Retin Eye Res.* 2006;25:397-424.
- Shibata M, Sugiyama T, Kurimoto T, et al. Involvement of glial cells in the autoregulation of optic nerve head blood flow in rabbits. *Invest Ophthalmol Vis Sci.* 2012;53:3726-3732.
- Metea MR, Newman EA. Glial cells dilate and constrict blood vessels: a mechanism of neurovascular coupling. *J Neurosci.* 2006;26:2862-2870.
- Song Y, Nagaoka T, Yoshioka T, Nakabayashi S, Tani T, Yoshida A. Role of glial cells in regulating retinal blood flow during flicker-induced hyperemia in cats. *Invest Ophthalmol Vis Sci.* 2015;56:7551-7559.
- Weinreb RN, Aung T, Medeiros FA. The pathophysiology and treatment of glaucoma: a review. *JAMA.* 2014;311:1901-1911.
- Antonetti DA, Barber AJ, Bronson SK, et al. Diabetic retinopathy: seeing beyond glucose-induced microvascular disease. *Diabetes.* 2006;55:2401-2411.
- Lott ME, Slocomb JE, Shivkumar V, et al. Comparison of retinal vasodilator and constrictor responses in type 2 diabetes. *Acta Ophthalmol.* 2012;90:e434-e441.
- Harris A, Anderson DR, Pillunat L, et al. Laser Doppler flowmetry measurement of changes in human optic nerve head blood flow in response to blood gas perturbations. *J Glaucoma.* 1996;5:258-265.
- Dallinger S, Dorner GT, Wenzel R, et al. Endothelin-1 contributes to hyperoxia-induced vasoconstriction in the human retina. *Invest Ophthalmol Vis Sci.* 2000;41:864-869.
- Zhu Y, Park TS, Gidday JM. Mechanisms of hyperoxia-induced reductions in retinal blood flow in newborn pig. *Exp Eye Res.* 1998;67:357-369.
- Takagi C, King GL, Takagi H, Lin YW, Clermont AC, Bursell SE. Endothelin-1 action via endothelin receptors is a primary mechanism modulating retinal circulatory response to hyperoxia. *Invest Ophthalmol Vis Sci.* 1996;37:2099-2109.
- Takagi C, Bursell SE, Lin YW, et al. Regulation of retinal hemodynamics in diabetic rats by increased expression and action of endothelin-1. *Invest Ophthalmol Vis Sci.* 1996;37:2504-2518.
- Polak K, Luksch A, Frank B, Jandrasits K, Polska E, Schmetterer L. Regulation of human retinal blood flow by endothelin-1. *Exp Eye Res.* 2003;76:633-640.
- Izumi N, Nagaoka T, Sato E, et al. Role of nitric oxide in regulation of retinal blood flow in response to hyperoxia in cats. *Invest Ophthalmol Vis Sci.* 2008;49:4595-4603.
- Ripodas A, de Juan JA, Roldan-Pallares M, et al. Localisation of endothelin-1 mRNA expression and immunoreactivity in the retina and optic nerve from human and porcine eye. Evidence for endothelin-1 expression in astrocytes. *Brain Res.* 2001;912:137-143.
- Ihara M, Noguchi K, Saeki T, et al. Biological profiles of highly potent novel endothelin antagonists selective for the ETA receptor. *Life Sci.* 1992;50:247-255.
- Eguchi S, Hirata Y, Ihara M, Yano M, Marumo F. A novel ETA antagonist (BQ-123) inhibits endothelin-1-induced phosphoinositide breakdown and DNA synthesis in rat vascular smooth muscle cells. *FEBS Lett.* 1992;302:243-246.
- Chakravarthy U, Gardiner TA, Anderson P, Archer DB, Trimble ER. The effect of endothelin 1 on the retinal microvascular pericyte. *Microvasc Res.* 1992;43:241-254.
- Meyer P, Flammer J, Luscher TF. Endothelium-dependent regulation of the ophthalmic microcirculation in the perfused

- porcine eye: role of nitric oxide and endothelins. *Invest Ophthalmol Vis Sci.* 1993;34:3614-3621.
21. Hein TW, Ren Y, Yuan Z, et al. Functional and molecular characterization of the endothelin system in retinal arterioles. *Invest Ophthalmol Vis Sci.* 2009;50:3329-3336.
 22. McMahon EG, Palomo MA, Moore WM, McDonald JF, Stern MK. Phosphoramidon blocks the pressor activity of porcine big endothelin-1-(1-39) in vivo and conversion of big endothelin-1-(1-39) to endothelin-1-(1-21) in vitro. *Proc Natl Acad Sci USA.* 1991;88:703-707.
 23. Ikegawa R, Matsumura Y, Tsukahara Y, Takaoka M, Morimoto S. Phosphoramidon inhibits the generation of endothelin-1 from exogenously applied big endothelin-1 in cultured vascular endothelial cells and smooth muscle cells. *FEBS Lett.* 1991; 293:45-48.
 24. Nagaoka T, Sakamoto T, Mori F, Sato E, Yoshida A. The effect of nitric oxide on retinal blood flow during hypoxia in cats. *Invest Ophthalmol Vis Sci.* 2002;43:3037-3044.
 25. Yoshida A, Feke GT, Mori F, et al. Reproducibility and clinical application of a newly developed stabilized retinal laser Doppler instrument. *Am J Ophthalmol.* 2003;135:356-361.
 26. Feke GT, Goger DG, Tagawa H, Delori FC. Laser Doppler technique for absolute measurement of blood speed in retinal vessels. *IEEE Trans Biomed Eng.* 1987;34:673-680.
 27. Nakabayashi S, Nagaoka T, Tani T, et al. Retinal arteriolar responses to acute severe elevation in systemic blood pressure in cats: role of endothelium-derived factors. *Exp Eye Res.* 2012;103:63-70.
 28. Riva CE, Cranstoun SD, Mann RM, Barnes GE. Local choroidal blood flow in the cat by laser Doppler flowmetry. *Invest Ophthalmol Vis Sci.* 1994;35:608-618.
 29. Ishikawa Y, Mine S. Amino adipic acid toxic effects on retinal glial cells. *Jpn J Ophthalmol.* 1983;27:107-118.
 30. Omae T, Nagaoka T, Tanano I, Yoshida A. Pioglitazone, a peroxisome proliferator-activated receptor-gamma agonist, induces dilation of isolated porcine retinal arterioles: role of nitric oxide and potassium channels. *Invest Ophthalmol Vis Sci.* 2011;52:6749-6756.
 31. Nagaoka T, Hein TW, Yoshida A, Kuo L. Simvastatin elicits dilation of isolated porcine retinal arterioles: role of nitric oxide and mevalonate-rho kinase pathways. *Invest Ophthalmol Vis Sci.* 2007;48:825-832.
 32. Hein TW, Xu W, Kuo L. Dilation of retinal arterioles in response to lactate: role of nitric oxide, guanylyl cyclase, and ATP-sensitive potassium channels. *Invest Ophthalmol Vis Sci.* 2006;47:693-699.
 33. Hein TW, Yuan Z, Rosa RH Jr, Kuo L. Requisite roles of A2A receptors, nitric oxide, and KATP channels in retinal arteriolar dilation in response to adenosine. *Invest Ophthalmol Vis Sci.* 2005;46:2113-2119.
 34. Kuo L, Davis MJ, Chilian WM. Endothelium-dependent, flow-induced dilation of isolated coronary arterioles. *Am J Physiol.* 1990;259:H1063-H1070.
 35. Rubanyi GM, Vanhoutte PM. Superoxide anions and hyperoxia inactivate endothelium-derived relaxing factor. *Am J Physiol.* 1986;250:H822-H827.
 36. Masaki T, Kimura S, Yanagisawa M, Goto K. Molecular and cellular mechanism of endothelin regulation. Implications for vascular function. *Circulation.* 1991;84:1457-1468.
 37. Moore D, Harris A, Wudunn D, Kheradiya N, Siesky B. Dysfunctional regulation of ocular blood flow: a risk factor for glaucoma? *Clin Ophthalmol.* 2008;2:849-861.
 38. Kohner EM, Patel V, Rassam SM. Role of blood flow and impaired autoregulation in the pathogenesis of diabetic retinopathy. *Diabetes.* 1995;44:603-607.
 39. Weinreb RN, Khaw PT. Primary open-angle glaucoma. *Lancet.* 2004;363:1711-1720.
 40. Lieth E, Barber AJ, Xu B, et al. Glial reactivity and impaired glutamate metabolism in short-term experimental diabetic retinopathy. Penn State Retina Research Group. *Diabetes.* 1998;47:815-820.
 41. Chihara E, Matsuoka T, Ogura Y, Matsumura M. Retinal nerve fiber layer defect as an early manifestation of diabetic retinopathy. *Ophthalmology.* 1993;100:1147-1151.
 42. Bloodworth JM Jr. Diabetic retinopathy. *Diabetes.* 1962;11:1-22.
 43. Bonaventure N, Roussel G, Wioland N. Effects of DL-alpha-amino adipic acid on Muller cells in frog and chicken retinae in vivo: relation to ERG b wave, ganglion cell discharge and tectal evoked potentials. *Neurosci Lett.* 1981;27:81-87.
 44. Pedersen OO, Karlsen RL. Destruction of Müller cells in the adult rat by intravitreal injection of D,L-alpha-amino adipic acid. An electron microscopic study. *Exp Eye Res.* 1979;28: 569-575.
 45. Yoshioka T, Nagaoka T, Song Y, Yokota H, Tani T, Yoshida A. Role of neuronal nitric oxide synthase in regulating retinal blood flow during flicker-induced hyperemia in cats. *Invest Ophthalmol Vis Sci.* 2015;56:3113-3120.